



BURNSIDE

Appendix B.1

Description of Soils Within the Region

(Source: Appendix C.1 – AESNP Transmission

System EIS, March 2001)

APPENDIX C.1

DESCRIPTION OF SOILS WITHIN THE REGION

Buganda Catena is characteristic of the Ntenjeru, Buikwe, Seeta and Nakisuga areas. It consists of shallow, skeletal soils developed from either quartzite or ironstone on summits and upper slopes and deep red or red-brown clay loams occurring on pediments. The latter are often associated with truncated and ferruginized soil profiles occurring in the lower sections of the pediments. Frequently, an extensive quartz dyke may be present in the hill summits with sandy or stony soils down slope.

The **Kyebe Catena** occurs in the areas of Ngogwe and south of Buikwe. The soils are closely related to the Buganda soils and particularly to the loamy associates of Buganda Catena. They differ from the latter in respect to relief that consists of hills with rounded summits and generally steeper and shorter pediments, and in short to medium grass savanna expanses of *Pennisetum purpureum*. Soils are lighter in texture (sandy loams) being probably derived from fine-grained sandstone and quartzites rather than schists as in the Buganda catena, and are generally shallower. The catena has a greater proportion of the area occupied by shallow, bouldery and excessively drained soils at the summits and steep upper slopes. Soils associated with strongly dissected remnants of the Buganda surface generally occur on gently rolling hills with rounded summits at 1,300 m to 1,500 m ASL. Most of these summits possess broken up and disintegrated boulders of laterite instead of extensive sheets as in the Buganda catena. The pediments are long and more gently inclined, with slope gradients of 5 to 8 percent and dissected by broad valleys.

The parent material of the upland soil series of these catenas is derived predominantly from the weathering products of Basement Complex gneisses and granitic rocks which give rise to red or brown loams with varying quantities of quartz gravel and stones.

The **Mirambi Catena** is common in the area of Lugazi, Nyenga and parts of Ngogwe. The Lugazi and Ngogwe areas are located close to the transmission line alignment. Analytically, the Mirambi soils are much less acidic and less leached than those of the Buganda and Kyebe catenas.

The soils of the **Mabira Catena** are generally ferralitic sandy clay loams with black waterlogged clays in the valley bottoms. The Mabira catena is characteristic of the entire Mabira Forest Reserve and the adjacent villages of Najjembe and Kawolo. The general relief of this series is somewhat steeper and the hills themselves are more ridge-like in appearance with long and generally rounded crests than the catenas previously described.

In this series the upland soil sequence is derived from two different bands of parent material. The summit and upper slopes are developed over relic ironstone and the pediment soils from weathering products of phyllite with minor occurrences of amphibolite, which give rise to red or yellow clays. The red associates of the Mabira Catena are strongly laterized and contain well developed murram horizons. The soil has a high clay content (well over 60 percent in the lower horizons) and generally a good nutrient status, particularly in respect of organic matter (4 to 6 percent carbon) and exchangeable bases.

Agriculturally, Mabira soil series are extensively cultivated and are generally very productive supporting all the commonly grown annual and perennial crops.

The Bujagali area is characterised by heavy loamy soils known as the Nakabango Catena that are generally rich in nutrients. These soils are usually between 0.15 m and 1.0 m deep. A variety of clays, light soils and sandy loamy soils are commonly found in valleys with a well-defined course and shallow alluvium in beds. There are also ferrisol (red) soils formed on basic rocks.

The Nakabango series is also characteristic of the sugarcane plantation area. It is associated with rolling to gently rolling hills with summits 1300 – 1500 m ASL. The general relief, climate and vegetation are very similar to that of the Mabira catena with which the Nakabango Series forms a complex at some points.

The parent material of the pediment soils is derived from weathering products of basic rocks, amphibolite schists and dolerites which on weathering give rise to bright red or reddish brown clays. It may easily be distinguished from all other upland soils in Buganda by its stickiness when wet and the relatively wide and irregular cracking on drying due to the almost complete absence of a quartz sand fraction.

The Nakabango medium soil (upper pediment) is normally well drained. The brown colour of this series may indicate impeded internal drainage; however, a low content of iron oxides, which have not been fully released from ferro-magnesian minerals, may also be a contributing factor.

The Nakabango soils are characteristically fertile and support a wide range of agricultural crops. The most productive coffee farms and the SCOUL sugar estate are situated on these soils.



Appendix B.2

Climatic Information for the Region

**(Source: Appendix C.2 – AESNP Transmission
System EIS, March 2001)**

APPENDIX C.2

CLIMATIC INFORMATION FOR THE REGION

Table 1: Mean Monthly Rainfall (R) and Potential Evapotranspiration (PET) in Different Parts of Mukono District (mm)

Month	Bbale		Ntenjeru		Nakifuma		Mukono		Buikwe		Buvuma	
	R	PET	R	PET	R	PET	R	PET	R	PET	R	PET
Jan	34	152	46	140	58	138	100	136	88	132	74	13
Feb	54	144	60	134	72	132	108	130	96	126	84	128
Mar	102	152	108	142	120	140	214	138	178	136	150	138
Apr	182	124	182	120	190	118	240	116	224	116	208	118
May	146	118	136	112	134	112	228	110	182	108	170	108
June	78	108	60	106	66	106	100	106	84	104	76	106
July	82	110	70	106	70	106	94	108	78	106	62	106
Aug	122	116	106	114	106	114	104	116	96	112	64	114
Sep	120	124	102	122	106	120	112	122	102	120	76	124
Oct	132	132	132	128	136	126	144	128	150	126	114	130
Nov	114	134	120	126	136	124	196	122	170	122	138	124
Dec	64	142	72	134	84	132	120	128	116	124	116	124
Yearly	1230	1556	1194	1484	1278	1468	1760	1460	1564	1432	1332	1454

Source: Meteorology Department, Kampala

Table 2: Meteorological Measurements at Kituza Agricultural Research Station

Month	Max Temp (°C)	Min. Temp. (°C)	Relative Humidity % (0600 hrs)	Relative Humidity % (1200 hrs)	Mean Sunshine Hrs/day	Mean Rainfall (mm)
Jan	27.5	14.5	88	64	7.1	68
Feb	27.2	15.1	92	66	6.0	108
Mar	26.8	15.8	89	72	5.5	175
Apr	26.1	16.0	89	74	5.1	203
May	25.8	15.9	89	75	5.6	165
Jun	25.6	14.9	88	71	6.0	83
Jul	25.1	14.3	91	71	5.3	62
Aug	25.7	14.2	90	71	5.2	81
Sept	26.4	14.7	86	69	5.8	130
Oct	26.8	15.5	86	69	4.9	160
Nov	26.7	15.2	86	70	5.5	243
Dec	26.5	14.7	88	70	6.9	132

Source: Meteorology Department, Kampala



BURNSIDE

Appendix B.3

**Report on Biodiversity of Proposed
Transmission Line**

**BIODIVERSITY OF
KEY SECTIONS OF THE
PROPOSED NEW BUJAGALI TO
KAMPALA TRANSMISSION LINE**

**With special reference to Mabira, Kifu and
Namyoya Central Forest Reserves, and the
Lubigi swamp**

Compiled and edited by Robert Kityo and Derek Pomeroy

Makerere University

September 2006

CONTENTS

<u>EXECUTIVE SUMMARY</u>	1
<u>1.0 GENERAL INTRODUCTION</u>	2
<u>1.1 SCOPE OF WORK</u>	3
<u>1.11 THE FOREST ECOSYSTEMS</u>	3
<u>1.12 THE WETLAND ECOSYSTEMS</u>	4
<u>1.2 FOREST SITES</u>	4
<u>1.3 WETLAND SITES</u>	5
<u>1.4 CONTRIBUTORS</u>	6
<u>2.0 VEGETATION</u>	7
<u>2.1 FOREST ECOSYSTEMS</u>	7
<u>2.11 METHODS</u>	7
<u>2.12 DATA ANALYSIS</u>	7
<u>2.2 RESULTS</u>	8
<u>2.21 MABIRA FOREST RESERVE</u>	8
<u>2.22 KIFU FOREST RESERVE (SITE 9, TOWER 66)</u>	13
<u>2.23 NAMYOYA FOREST RESERVE (SITE 10, TOWER 51)</u>	14
<u>2.24 SIMILARITIES AMONG MABIRA, KIFU AND NAMYOYA FOREST RESERVES</u>	14
<u>2.25 GENERAL DISCUSSION AND COMPARATIVE OVERVIEW WITH PREVIOUS STUDY</u>	15
<u>2.26 SPECIES OF CONSERVATION CONCERN AND ECONOMIC IMPORTANCE</u>	16
<u>2.27 ALIEN INVASIVE SPECIES: BROUSSONETIA PAPYRIFERA, TITHONIA DIVERSIFOLIA AND LANTANA CAMARA</u>	16
<u>2.3 WETLAND ECOSYSTEMS – LUBIGI SWAMP</u>	17
<u>2.31 INTRODUCTION</u>	17
<u>2.32 METHODS</u>	18
<u>2.33 RESULTS</u>	18
<u>2.34 SIMILARITY AMONG SITES</u>	18
<u>3.0 INVERTEBRATES: BUTTERFLIES AND DRAGONFLIES</u>	20
<u>3.1 INTRODUCTION</u>	20
<u>3.2 METHODS</u>	20
<u>3.3 RESULTS: DRAGONFLIES</u>	20
<u>3.4 RESULTS: BUTTERFLIES</u>	21
<u>3.5 DISCUSSION</u>	22
<u>4.0 AMPHIBIANS AND REPTILES</u>	23
<u>4.1 INTRODUCTION</u>	23
<u>4.2 OBJECTIVES</u>	23
<u>4.3 METHODS</u>	24
<u>4.4 RESULTS</u>	25
<u>4.41 AMPHIBIANS</u>	25
<u>4.42 REPTILES</u>	27
<u>4.43 SPECIES OF CONSERVATION CONCERN</u>	28
<u>4.5 DISCUSSION</u>	28
<u>5 BIRDS</u>	29
<u>5.1 FOREST BIRDS: INTRODUCTION</u>	29
<u>5.2 METHODS</u>	29
<u>5.3 RESULTS AND DISCUSSION</u>	30
<u>5.4 SWAMP BIRDS</u>	31
<u>6.0 MAMMALS</u>	33
<u>6.1 INTRODUCTION</u>	33

<u>6.2</u>	<u>METHODS</u>	33
<u>6.3</u>	<u>RESULTS</u>	33
<u>6.31</u>	<u>MABIRA FOREST</u>	33
<u>6.32</u>	<u>KIFU AND NAMYOYA FORESTS</u>	35
<u>6.34</u>	<u>IMPORTANCE OF THE FOREST ALONG THE TRANSMISSION LINE FOR MAMMALS</u> ..	36
<u>6.4</u>	<u>LUBIGI WETLANDS</u>	37
<u>7.0</u>	<u>SUMMARY OF IMPACTS AND PROPOSED MITIGATION MEASURES</u>	39
<u>7.1</u>	<u>LIKELY IMPACTS</u>	39
<u>7.2</u>	<u>OVERALL IMPACT ASSESSMENT</u>	40
<u>7.3</u>	<u>MANAGEMENT OPTIONS AND MITIGATION MEASURES</u>	41
	<u>REFERENCES</u>	43

LIST OF TABLES

<u>Table P1: Number of species recorded from the Forests Reserves surveyed in the different families of plants.....</u>	9
<u>Table P2: Diameter class distributions and stem densities for the study sites in Mabira forest.....</u>	10
<u>Table A1. Dragonfly species abundance categorization.....</u>	21
<u>Table A2. Summary of species recorded by this study.....</u>	21
<u>Table A3. Summary of butterfly species with their respective ecological types according to Davenport et al (1996).....</u>	22
<u>Table A4. Butterfly species abundance data.....</u>	22
<u>Table H1: Amphibian encountered in Mabira Forest Reserve.....</u>	26
<u>Table H2: Amphibian encountered in Lubiji Swamp.....</u>	26
<u>Table H3: Reptilian fauna recorded from Mabira & Kifu Forests.....</u>	27
<u>Table H4: Reptilian fauna recorded from Lubiji Swamp.....</u>	27
<u>Table B1. Bird descriptors. No globally-threatened species were observed. The species' preferred habitats and migratory status are also indicated in Table B4 and Appendix B1.....</u>	29
<u>Table B2. Summary of numbers of 'tree birds' and other species recorded in Mabira Forest in 2001 and 2006. There were 16 counts in 2001, and 8 in 2006.....</u>	30
<u>Table B3. Summary of 'tree birds' and other species recorded in Kifu Forest in 2001 and 2006. In 2001, two TSC counts were made, with three in 2006.....</u>	30
<u>Table B4. Summary of 'tree birds' and other species recorded in Namyoya Forest in 2001 and 2006. Two counts were made in 2001, but only one in 2006.....</u>	30
<u>Table B5. Globally and regionally threatened species recorded in Mabira Forest in 2006.....</u>	31
<u>Table B6. Summary of bird data from the six swamp sites (as listed in Appendix G2). The categories of bird types are listed in Table B4.....</u>	32
<u>Table M1 Trap and netting success recorded at the different sampling locations.....</u>	34
<u>Table M2 Proportionate record of the known mammalian richness in Mabira FR in 2001 and 2006.....</u>	34
<u>Table M3. Mammal species recorded from Kifu and Namyoya Central Forest Reserves.....</u>	36
<u>Table M4. Mammal records for Lubiqi Wetlands.....</u>	38
<u>Table 7.1. Potential impacts in the forested areas.....</u>	39
<u>Table 7.2. Potential impacts in the wetland sites.....</u>	40

LIST OF FIGURES

<u>Figure P1: Species-area curves for the combined studied sites in Mabira forest reserve.....</u>	10
<u>Figure P2: Mabira forest tree dbh distributions for study sites 1 – 8.....</u>	11
<u>Figure P3: Cluster analysis of 8 sites of Mabira forest based upon the presence or absence of 275 plant species. The cut-off for the classification of sites is the Simple Matching Coefficient (SM) of 0.31, represented by the dotted line.....</u>	12
<u>Figure P4: Diameter size classes for the trees found on the southern side of the existing powerline in Kifu forest reserve.....</u>	14
<u>Figure P5: Cluster analysis of the three forest reserves based upon the presence or absence of 315 plant species using a Simple Matching coefficient (SM) cut-off of 0.2.....</u>	15
<u>Figure P6: Cluster analysis of the six sample sites in Lubiqi swamp based upon the presence or absence of 124 plant species. Site 1 (Kazinga 1), Site 2 (Kazinga 2), Site 3 (Lubanyi 1), Site 4 (Lubanyi 2), Site 5 (Nakawudde) and Site 6 (Nganda-Nansana).....</u>	19
<u>Figure M1. The cumulative number of species of small mammals (rodents & insectivores) recorded overall along the sample transects.....</u>	35

APPENDICES: GPS locations and species lists

<u>Appendix F1. Figures (Maps).....</u>	46
<u>Appendix G1. GPS locations of landmarks that were noted in Mabira, Kifu and Namyoya Forests.....</u>	52
<u>Appendix G2. GPS locations of landmarks that were noted along the western border of Lubiqi Swamp.....</u>	53

<u>Appendix P 1: Plant species recorded from Mabira, Kifu and Namyoya forest reserves</u>	54
<u>Appendix P2: Plant species list for Lubigi swamp</u>	64
<u>Appendix A1 Dragonfly species list</u>	67
<u>Appendix A2. Butterfly species list</u>	68
<u>Appendix A3. Comparison of butterfly species recorded from Mabira forest.</u>	74
<u>Appendix B1. Summary of bird count data from Mabira forest. The columns T1-8 and M1-8 are the TSC scores, and numbers of birds mist-netted, respectively. Mean values are also given</u>	77
<u>Appendix B2. Summary of bird count data from the smaller forests, and from the wayleave in Mabira, which is a non-forest site. There were two TSCs in the Mabira wayleave area (T9, 10); three in Kifu (K1, 2,3 and the mean KM) and one at Namyoya (N)</u>	80
<u>Appendix B3. Bird records from Lubigi wetland</u>	83
<u>Appendix M1. Mammal species recorded along the various transects and locations surveyed in Mabira Forest.</u>	86
<u>Photoplates</u>	87

EXECUTIVE SUMMARY

1. The proposed route of the Bujagali–Kampala transmission line passes through three Central Forest Reserves (CFRs: Mabira, Kifu and Namyoya) and along the edge of a major wetland (Lubigi). In each of these we conducted surveys to assess their biological importance, which included biodiversity, rare or endangered species and important ecosystems. To do this, we surveyed plants, dragonflies, butterflies, frogs, reptiles, birds and mammals at each site. All members of the team are experienced in such work.
2. These surveys repeated and extended those made in the three CFRs in 1999. No significant differences were found, nor are any endangered species likely to be adversely affected by the new line. Nevertheless, Mabira is the only major forest along the northern shores of Lake Victoria and it supports a high biodiversity, including a large number of species of conservation concern. Consequently, it is important to mitigate the forest loss, whether by improving some degraded parts of Mabira, or by reforestation elsewhere. In either case, it is of course the natural forest which has the most value. It is also important to prevent the fragmentation of the forest preventing species from moving from the southern to the northern blocks, and vice-versa. We therefore propose valley corridors to retain a link between them. The least distributed forest is to the north of the existing wayleave so that, if a way can be found to do so, the new line should be to the south of the existing line.
3. The two smaller forests have already been extensively changed, and such parts as are still forest are almost entirely planted with exotic species, notably eucalyptus. Their biodiversity value is low.
4. Lubigi swamp, despite its proximity to Kampala, has not been extensively altered and its biodiversity value is still high, and we recommend that it should be formally protected. The erection of pylons along its north-western border will not greatly affect the swamp, particularly as a major highway is taking a much longer part of it.
5. Overall, the construction of the proposed transmission line will not have a very serious impact on the biological value of the areas affected: and these impacts can be mitigated fairly easily.
6. The substantial amounts of data collected have been deposited in the National Biodiversity Data Bank in the Makerere University Institute of Environment and Natural Resources and they will provide a valuable basis of comparison for future studies.

1.0 GENERAL INTRODUCTION

Forest Sites

Mabira is the largest Forest Reserve in Central Uganda (Davenport *et al* 1996), found in an area of gently undulating land interrupted by flat-topped hills that are remnants of the ancient African peneplain (Howard 1991). During the cooler, drier parts of the Pleistocene Period, there was no forest in this area. The forests around the northern shores of Lake Victoria originated only 10-12,000 years ago, as the climate became warmer and wetter (Hamilton, 1982).

In a review of the biodiversity importance of 65 of Uganda's forests, the then Forest Department (2002) ranked Mabira 24th (p145), although somewhat higher in terms of rare species. (Neither Namyonya nor Kifu were assessed). The proposed zonation of the forest included a Nature Reserve of 73 km², whose southern limits would be some 5 km north of the existing transmission line.

The vegetation of Mabira is classified as medium-altitude moist semi-deciduous forest (Langdale-Brown *et al*, 1964) and is considered to be of sub-climax stage, highly disturbed and heavily influenced by man. The forest, which covers an area of 306 km² is found 54 km east of Kampala and only 26 km west of Jinja. Such proximity to large urban centers and also being located in areas of fairly high human populations, puts considerable strains onto the forest for forest products.

Both the National Forest Authority (NFA) and private owners have replanted a large section of Kifu Central Forest Reserve (CFR) with plantation trees. The southern side of the existing powerline was replanted with *Maesopsis eminii* but other indigenous species still exist, scattered among the *Maesopsis eminii* trees. The common trees include *Funtumia elastica*, *Alchornea cordifolia*, *Antiaris toxicaria* and *Celtis mildbraedii*. Shrubs and herbaceous species such as *Glycine wightii*, *Marantochloa leucantha*, *Pteris dentata*, *Renealmia congolana*, *Acanthus pubescens*, *Acalypha bipartita*, *Acalypha ornata*, *Scutia myrtina* and *Rubus apetalus* characterize the understorey. The northern side has been replanted with *Araucaria* spp. after clearing the natural vegetation of the area except for the herbaceous vegetation that is regularly weeded out to enable proper growth of the *Araucaria*.

Namyoya CFR. is predominantly *Eucalyptus* woodlots on the northern side of the existing powerline but with a few indigenous tree species still surviving, including for example *Funtumia elastica*, *Alchornea cordifolia*, *Sapium ellipticum* and *Erythrina abyssinica*. The southern side of the powerline at the point we conducted the surveys, has gardens and harvested *Eucalyptus* woodlots. Some of the herbaceous plants and shrubs recorded are characteristic of disturbed areas and include *Acanthus pubescens*, *Ageratum conyzoides*, *Bidens pilosa*, *Conyza floribunda* and *Vernonia amygdalina*.

The surveys we are documenting in this report were planned to update the terrestrial ecological assessments of the Mabira, Namyoya (Mwola) and Kifu forest reserves which were included in the 2001 Transmission System Environment Impact Statement (EIS). Consequently, most of the data were collected from the same places, and using similar methods, to the earlier survey, although with additional taxa, as described in Section 1.1.

Wetland sites

Wetlands have distinct vegetation characteristics and in addition the soils found under a wetland are quite different from normal soils because they are formed under low oxygen conditions and are very heavy with clay or have large amounts of plant remains known as peat.

We have not found very much in the way of documentation about Lubigi wetlands except the brief profile of Namakambo (2000). According to Namakambo (2000), Kampala district has numerous wetlands that are closely linked with the topography and drainage of the district. Although very near to Kampala city and Kampala administrative district, Lubigi swamp is currently under the jurisdiction of Wakiso district. Lubigi wetlands are among the largest wetlands in the district, not very far removed from its border with Kampala district. It is permanently waterlogged, being fed by rivers Mayanja, Bwaise, Nabisasiro, Nalukolongo and Kiwunya.

Lubigi wetlands extend through Lubaga and Kawempe subcounties and are located approximately 7.5 km west of Kampala city (Namakambo 2000). The same author indicated that the wetlands could be accessed at several locations on Masaka, Hoima, Mityana and Sentema roads. Several villages, including Busega, Natete, Bulenga, Nakuwadde, Lubanyi, Masanafu, Kawala, Nganda and Namungoona surround the wetlands. All these are suburban villages, which are densely populated with resultant impacts on the condition of the wetland near them.

Namakambo (2000) described the swamp as being dominated by papyrus with patches of *Loudetia* sp., *Typha* sp., *Miscanthedium* sp., *Echinochloa* sp., and *Phoenix* palms. During our surveys we have also found communities of these plants to still be characteristic of the area (Plate 11). At the time of writing the author indicated that the areas adjacent to the wetland were cultivated except the immediate fringe. The only other wildlife that this author mentions is the presence of Sitatunga. At the time Namakambo wrote, the Lubigi swamp had no conservation status and no conservation measures had been proposed. Our consultations with the Wetlands Inspectorate Division revealed that the Lubigi wetland has no conservation status in any part of it. At the present time, the new Northern bypass highway, currently under construction traverses the eastern length of Lubigi wetlands. The road project has resulted in infilling in those sections of the swamp crossed by the highway. The EIA report on that project (2001) does recognize the ecological significance of the swamp, and recommend, the enforcement of wetland regulations.

Wetlands were not included in the 2001 transmission line EIS report. We selected six sample sites following a survey along the most accessible parts of the swamp. The locations were selected were chosen because they represented a fair sample of the available microhabitats along the edge of the swamp.

1.1 SCOPE OF WORK

We have conducted surveys to provide answers to question posed by our terms of reference, defined in the scope of work.

1.11 THE FOREST ECOSYSTEMS

1. Transect surveys of plant species at sites used for the 2001 EIS, including species presence/absence, size (dbh), and accumulation curves for plant species at each transect site.
2. Plant stock inventories for conducting stock assessments.

3. Inventories of birds at each of the same transects used for the 2001 EIS.
4. Inventories of terrestrial vertebrates and two groups of invertebrates and associated habitats at each of the transects.
5. Assessment of affected habitats and species including:
 - (i) Identification of any Critically Endangered or Endangered species as defined in the IUCN Red list of threatened species.
 - (ii) Identification of any critical Habitat as defined in the International Finance Corporation's Performance standard 6 (Biodiversity Conservation and Sustainable Natural Resource Management).
6. Assess impacts of the proposed transmission line construction on the features as identified in 1 – 5 above, and suggest appropriate measures for avoidance and mitigation of adverse effects, which can be incorporated into the SEAs Environment Action Plan (EAP). Such measures would as far as is practicable aim to integrate with any measures that are formulated to mitigate effects on tourism activities, the Mabira CFR management plan, and/or enhance the Kalagala offset plan.
7. Identify and make recommendations for capacity building or community-based natural resource management initiatives.

1.12 THE WETLAND ECOSYSTEMS

We were also required to conduct surveys to enable us to evaluate the ecological values of Lubigi wetlands. Among others, this task required us to: -

1. Consult with the Wetland Inspectorate Division, to ascertain the conservation status of the wetland, obtain and review available data on ecological conditions of the wetland.
2. Conduct detailed surveys of plants, birds, mammals, reptiles, amphibians and invertebrates in six sampling locations along the edge of the wetland.
3. Survey the full length of the edge of the swamp to identify and map the distribution of different ecological communities of the wetland and to use these to produce an ecological communities map detailing their distribution along the western edge of the swamp that will be traversed by the transmission line.
4. Prepare a description of the different ecological communities pointing out the defining characteristics of each.
5. Identify species of economic and conservation importance, including:
 - (i) Identification of any critically endangered or endangered species as defined in the IUCN Red list of threatened species.
 - (ii) Identification of any critical Habitat as defined in the International Finance Corporation's Performance standard 6 (Biodiversity Conservation and Sustainable Natural Resource Management).
6. Assess impacts of the proposed transmission line construction on the future identified in 1 – 5 above and suggest appropriate measures for avoidance and mitigation of adverse effects, which can be incorporated into the SEAs Environment Action Plan (EAP). Such measures should as far as is practicable aim to integrate any community development measures into the EAP.

1.2 FOREST SITES

In the forests (Mabira, Kifu & Namyoya), we revisited the same 10 locations that we used for the 2001 report. Eight transects were located in Mabira, and one each in Kifu and Namyoya

(Appendix G1).

The surveys in Mabira forest were conducted in the area of the forest bounded to the east by pylon 179 and pylon 144 to the west (coordinates are in Appendix G1). This stretch of forest covers a little over 7 km. The stretch of forest between Pylon 179 up to about Pylon 172 covering a distance of a little over 1 km, is largely characterized by paper mulberry. From Pylon 172 until 140 a little before Wasswa village, most of the way except for a short stretch between Pylon 159 & 158, the transmission line runs through more or less secondary forest with a narrow fringe of paper mulberry at the edge.

Starting a short distance after Pylon 140 (N00°26.3183', E032°59.3487'), and continuing westwards, the wayleave is under cultivation growing a variety of crops, including cassava, potatoes, yams, and bananas among others (Plates 3a-d).

The botanical surveys in Kifu CFR were done along a transect at pylon 166, for the other taxa (butterflies, birds and mammals) we moved around to as far as pylon 164. The forest to the north of the transmission line is now a plantation forest growing largely *Auricularia cunninghamii* while that to the south is largely planted with *Maesopsis eminii* but is covered in dense bushes of secondary growth.

Namyoya CFR, as far as could be observed, has been planted with *Eucalyptus*, which in the younger trees is intercropped with subsistence crops including maize, cassava, yams and potatoes. A small stand of trees measuring about 195 x 50 meters survives on the northern side of the transmission line in this area opposite pylon 51 as the only natural enclave. In this area surveys were conducted around the areas adjacent to Pylon 51 and were not restricted to a transect as was the case in Mabira and Kifu Forests.

Appendix G1 presents GPS recordings for those locations in Mabira, Kifu and Namyoya from which or around which data of one sort or another were taken. The GPS units were set for:

- 1 Map Datum WGS 84,
- 2 Units – Metric
- 3 North Ref – Mag.

The pylon GPS readings represent the locations of the transects and all these were taken from the wayleave as far away from the existing transmission lines as we could get without the canopy cover influencing how many satellites the GPS could receive.

Many waterways, potential waterways and/or points were dry at the time we conducted these surveys except for River Waliga in the area of site 3 and in six other locations west of site 5. We made note of these in particular because it seemed they would be important for provision of water for amphibians as well as other wildlife in Mabira to drink for most of the year and as potential points for maintaining corridors between the northern and southern parts of the forest for understorey species. Most of the amphibian records from Mabira were also from these points that had water at the time of these surveys.

The landmarks labeled “stream with water” for Mabira forest (Appendix GI) represent the only other locations that had either standing or running water at the time of the present surveys. The landmark temporary pond had a muddy puddle remaining at the edges of which spoors of two carnivores, the marsh mongoose and serval cat were recorded. All other landmarks had different notable features as are implied in the landmark descriptor.

Field work was conducted at these sites between 30 June and 7 July 2006.

1.3 WETLAND SITES

We conducted a survey along the accessible edge of Lubigi swamp to describe and map the different ecological communities along the edge of the swamp. We also used this exercise to identify the most suitable and representative locations for conducting the detailed biodiversity surveys.

The GPS units were set as described for Section 1.2 for the forest sites. Appendix G2 contains details of the locations that were visited and described for purposes of mapping the ecological communities around the swamp. Six of these locations (sites 1 – 6) were then used for the detailed ecological surveys.

Additional points which were not necessarily along the edge of the swamp to be traversed by the transmission line, were also recorded because they represented areas where the swamp was already being impacted by human activities such as brick making, dumping refuse, burning bones, mining for sand and clay, papyrus harvesting or cultivation.

Field work in the wetlands was conducted between 10 and 14 July 2006.

1.4 CONTRIBUTORS

The following people undertook the field work, whilst the preparation of relevant sections of this report was undertaken by those in bold.

Plants	Paul Ssegawa
Invertebrates	Perpetra Akite and Andrew Ochama
Amphibians and reptiles	Stephen Kigoolo
Birds	Eric Sande, Achilles Byaruhanga and Derek Pomeroy
Mammals	Robert Kityo and Rebecca Nalunkuma
Robert Kityo also co-ordinated the field work.	

2.0 VEGETATION

2.1 FOREST ECOSYSTEMS

2.11 METHODS

The same study sites were used as in the previous survey, which was carried out in 1999 and reported in AESNP (2001). Appendix G1 gives the locations at which transects of 500 m were established.

Along each transect, plots of 0.02 ha. (20 x 10 m) were used. These were located alternately along a transect at a spacing of 50 m. Each transect had ten quadrats located on the northern and southern sides of the existing power line. The transects ran in a north-south direction. Using this approach, a total of 80 plots were sampled at the eight sites located in Mabira forest. Poles and flagging tape were used to mark the plots. Ground distance was measured using a 50-m tape without correcting for the slope. Each woody plant (tree, shrub or climber) of diameter ≤ 3 cm encountered in the plots was identified, enumerated and its diameter measured at 1.3 m (diameter at breast height, dbh) using a 5-m diameter tape. The following decisions on which plants to include in the plot were taken: plants branching below 1.3 m, had their individual stem diameters measured and averaged. For trees with large diameters and buttresses extending outside the plot boundaries, only those individuals with the mid point of the base inside the plot were included. Plants whose stems grew into the plot but with their bases outside were not included. This was particularly the case for climbers. Plots of 10 m x 10 m were used to census plants in the wayleave at each study site. Collections were made of plants that could not be identified in the field were brought to the Makerere University Herbarium for identification. Identification was done with the help of taxonomic literature such as Flora of Tropical East Africa (FTEA), and Katende *et al* (1995).

2.12 DATA ANALYSIS

Site similarity

A compilation of the species at site level was done indicating the individual abundances at different sites using the DAFOR scale. The number of times a species appeared in the quadrats gave a good indication of its abundance in a particular study area.

Cluster analysis (Kovach, 1999) was used to group sites according to their respective species assemblage compositions to determine the degree of similarity among them. This was based on the presence/absence data. Cluster analysis is a technique that sorts objects (such as sampling units) into groups or clusters based upon their overall resemblance to one another (Ludwig & Reynolds, 1988). Dendrograms were developed for the sites within Mabira forest and also for comparison of the three forests (i.e. Mabira, Kifu and Namyoya) based on the presence/absence of species.

Species richness estimations

Because observed species (Sobs) as a species richness estimator, underestimates the number of species in any given homogeneous area, statistical methods have been developed to reduce this bias (see Colwell & Coddington, 1994). For tropical tree populations, Chazdon (1998) considered

that non-parametric extrapolation methods are the most efficient for predicting population richness from samples. The main non-parametric estimators are the first- and second-order Jackknife (Heltsh & Forrester, 1983). Jackknife 1 (Jack 1) and Jackknife 2 (Jack 2) species richness estimators (Colwell & Coddington, 1994) were used to estimate the species richness. Jackknife 1 and Jackknife 2 are incidence-based non-parametric estimators that were developed to estimate the number of species in a random sample from a single population (Colwell & Coddington, 1994; Guralnick & Van Cleve, 2005). Incidence-based estimators use the relative rarity and commonness of species in subsamples (plots) of the complete sample to estimate richness. Species area curves were also plotted to determine the adequacy of the sampling effort (Bhatt & Sanjit, 2005). Species richness estimations were calculated using the ESTIMATES 6b1a program (Colwell, 2001).

2.2 RESULTS

2.21 MABIRA FOREST RESERVE

A total of 274 species belonging to 242 genera and 76 families were recorded. For the trees, a total of 94 species with 1374 individuals were recorded with a combined basal area of 52.3 m² in eighty 0.02 ha. plots. A total of 34 families were represented by one species each, and the others were represented by 2 – 25 species each (Table P1 and Appendix P1). About 48% of the species were categorised as rare because they were recorded in one out of eight sites.

The family Fabaceae had the highest number of species (25) followed by Poaceae and Euphorbiaceae with 18 and 17 respectively. The most represented genus was *Ficus* with 9 species. The commonest species included *Dracaena fragrans*, *Acalypha neptunica*, *Argomuelleria macrophylla* and *Broussonetia papyrifera*. These were recorded in all eight sites. The total number of trees recorded represented 34.1% of the total species recorded whereas herbs, climbers, grasses and shrubs represented 22.9%, 19.3%, 6.9% and 16.4% respectively. The highest observed number of species was recorded at Site 3 (Towers 167–170) and represented 38.2% of the total species recorded during the Mabira forest survey. The least number of observed species were recorded at Sites 1 (Tower 179) and 2 (Tower 174) representing 25.8% and 26.5% respectively (see Appendix P1).

The estimated total number of species by first-order Jackknife and second-order Jackknife species richness estimators for the eight sites sampled was 419 and 523 species respectively. The shapes of the species-area curves (Figure P1) indicate that continued sampling would yield more species. This is further reinforced by the relatively high estimated species richness, which indicates the presence of many rare species (Figure P1).

Forest stand density and basal area

The eight Mabira forest sites sampled had a stem density of 1374 in 1.6 ha (mean density 859 stems ha⁻¹). Stem density was greatest (1315 stems per ha) in Site 1 and least (630 stems per ha) in Site 7. Stem density was more or less similar in Sites 1 and 2. Basal area was highest in Site 3 representing 20% of basal area per ha for all the study sites while the least was recorded in Site 8 representing 3.9% (see Table P2).

Table P1: Number of species recorded from the Forests Reserves surveyed in the different families of plants

Family	Mabira	Kifu	Namyoya	Family	Mabira	Kifu	Namyoya
Acanthaceae	9	4	1	Marantaceae	1	1	0
Alangiaceae	1	0	0	Melastomataceae	1	0	0
Amaranthaceae	4	1	0	Meliaceae	8	0	0
Anacardiaceae	1	1	0	Menispermaceae	2	0	1
Annonaceae	2	0	0	Moraceae	15	4	4
Apocynaceae	8	2	1	Musaceae	1	0	0
Araceae	2	1	0	Myrsinaceae	2	1	0
Aristolochiaceae	2	0	0	Myrtaceae	3	0	2
Asclepiadaceae	3	1	0	Nyctaginaceae	1	0	0
Aspleniaceae	1	0	0	Ochanaceae	1	0	0
Asteraceae	9	5	6	Oleaceae	1	0	0
Auracariaceae	0	1	0	Oleandraceae	1	0	0
Balanitaceae	1	0	0	Oxalidaceae	1	0	0
Bignoniaceae	3	1	0	Passifloraceae	3	0	0
Burseraceae	1	0	0	Phytolaccaceae	1	0	0
Capparidaceae	4	0	0	Piperaceae	1	1	0
Cecropiaceae	1	0	0	Plumbaginaceae	1	0	0
Celastraceae	1	1	0	Poaceae	18	8	5
Combretaceae	1	0	0	Pteridaceae	1	1	0
Commelinaceae	4	2	0	Rhamnaceae	3	3	0
Connaraceae	4	0	0	Rosaceae	1	1	0
Convolvulaceae	4	4	2	Rubiaceae	7	2	0
Cucurbitaceae	3	1	0	Rutaceae	9	4	0
Cyperaceae	2	2	2	Sapindaceae	13	3	0
Davalliaceae	2	0	0	Sapotaceae	6	0	0
Dichapetalaceae	1	0	0	Simaroubaceae	1	0	0
Dilleniaceae	1	0	0	Smilacaceae	1	1	0
Dioscoreaceae	1	0	0	Solanaceae	5	1	1
Dracaenaceae	2	0	0	Sterculiaceae	5	0	0
Ebenaceae	1	0	0	Thelypteridaceae	1	1	0
Euphorbiaceae	17	10	6	Thymelaeaceae	0	1	0
Fabaceae	25	10	5	Tiliaceae	3	0	0
Flacourtiaceae	2	0	0	Ulmaceae	9	2	0
Guttiferae	0	1	1	Umbelliferae	1	0	1
Hernandiaceae	1	0	0	Urticaceae	1	0	0
Labiatae	3	0	1	Verbenaceae	6	1	1
Leeaceae	1	1	0	Violaceae	1	0	0
Malpighiaceae	1	0	0	Vitaceae	2	0	0
Malvaceae	5	4	3	Zingiberaceae	4	1	0




Figure P1: Species-area curves for the combined studied sites in Mabira forest reserve.

Table P2: Diameter class distributions and stem densities for the study sites in Mabira forest.

dbh (cm)	Species richness	Stem density	Number of stems at each site							
			1	2	3	4	5	6	7	8
3 – 10	80	653	175	115	73	70	65	62	46	47
11 – 20	53	394	61	68	34	55	39	45	49	43
21 – 30	43	168	25	26	14	21	21	24	15	22
31 – 40	33	65	2	5	3	11	17	11	5	11
41 – 50	25	44	0	3	3	9	12	5	6	6
51 – 60	9	14	0	2	0	0	5	2	3	2
Over 60	19	36	0	2	9	4	4	7	2	8
Totals			263	221	136	170	163	156	126	139
Stem density per ha			1315	1105	680	850	815	780	630	695
Basal area (m² ha⁻¹)			17.3	25.3	52.4	36.4	50.8	44.3	24.8	11.4

Forest stand structure

The population structure of the forest study site stands was reverse J-shaped with dbh frequency and basal area distribution in various size classes having a similar pattern in all the sites except for the largest (i.e. dbh over 60 cm) size class, which was greatest in Site 3 and absent in Site 1 and intermediate in Sites 4 and 5 (Figure P2). Tree species richness and density consistently decreased with increasing stem size classes except in the last class (i.e. Over 60 cm, see Table P2). The lowest size class captured 29% of the species richness, 47.5% of the forest stem density, and there was about a 9-fold decrease in species richness and a 47-fold decrease in stem density in the 51 – 60 cm dbh class.




Figure P2: Mabira forest tree dbh distributions for study sites 1 – 8.

Site similarity

Species presence or absence was scored in the 8 sites and provided the basis for cluster analysis, which provides evidence of likeness of species assemblages among the 8 study sites (Figure P3).




Figure P3: Cluster analysis of 8 sites of Mabira forest based upon the presence or absence of 275 plant species. The cut-off for the classification of sites is the Simple Matching Coefficient (SM) of 0.31, represented by the dotted line.

Sites 5 and 6 had the highest Simple Matching coefficient (SM) of 0.789 followed by Sites 7 and 8 clustering with 0.771 while Sites 1 and 2 clustered at 0.681. Other sites clustered at a lower value, and using a minimum SM of 0.31 for defining clusters (i.e. the dotted line in Figure P3), the analysis produced three distinct groups of sites M1, M2 and M3 (Figure P3).

The cluster analysis in Figure P3 shows that Site pairs 5-6 and 7-8 are similar though not closely, in terms of presence or absence of species. In cluster M2, Site pair 3 and 4 has a lower similarity compared to the site pair 1 and 2. However, in general terms, there is a recognizable east-west gradient in the sites clustering, probably due to the noticeable decrease in levels of disturbance, thus supporting particular species assemblages, as one moves from east to west.

Cluster analysis produced three distinct groups; Cluster M1 consisted of those sites on the western side of Mabira that have experienced less disturbances particularly to the northern side of the existing power transmission line, but which are also less species rich overall (see discussion of M2 cluster below). These sites represent larger diameter size classes with sparse undergrowth on the northern side. They represent part of the buffer zone of Mabira forest. However, the climbers, *Agelaea ugandensis*, *Acacia pentagona* and *Teramnus labialis* in the undergrowth and fewer large diameter trees dominate the southern plots. *Acalypha neptunica* is also a common understorey shrub. The characteristic tree species in these sites include *Albizia gummifera*, *Albizia glaberrima*, *Chrysophyllum albidum*, *Celtis mildbraedii*, *Alstonia boonei*, *Trilepisium madagascariensis* and *Pseudospondias microcarpa*. The southern side had fewer individuals of trees and of lower diameter classes with bigger canopy gaps and dense climber tangles. Commoner trees included *Antiaris toxicaria*, *Celtis mildbraedii* and *Funtumia elastica*. *Solanum mauritianum*, *Acalypha neptunica*, *Draceana fragrans* and the climber *Acacia pentagona* mainly constituted the shrub layer.

Cluster M2 sites are mainly influenced by the riverine conditions along transect 3. This cluster has the most species rich sites and there are characterised by intermediate diameter class trees. The characteristic tree species include *Celtis durandii*, *Aphania senegalensis*, *Teclea nobilis*, *Chrysophyllum albidum*, *Ficus polita* and *Blighia unijugata*. The herbs, *Pseuderanthemum ludovicianum* and *Pollia condosata* dominate the forest floor. Cluster M3 consists of sites 1 and 2, which are dominated by *Broussonetia papyrifera*. These are sites that were previously heavily encroached until 1992 when the encroachers were evicted. *Broussonetia papyrifera* prefers open areas that will enhance its regeneration. This, coupled with its invasiveness has enabled it to proliferate in these two sites. However, a few individuals of *Albizia grandibracteata*, *Celtis mildbraedii* and *Celtis wightii*, with the shrubs, *Acalypha neptunica* and *Argomuellera macrophylla* are struggling underneath the *Broussonetia papyrifera*. This cluster of sites registered the lowest species richness (Appendix P1).

Herbaceous plants that include *Lantana camara*, *Tithonia diversifolia*, *Lantana triphylla*, *Cynodon dactylon*, *Indigofera spicata* and *Sida rhombifolia* dominate the wayleave. It is regularly used to graze cattle and these species can withstand both the grazing and maintenance pressure.

2.22 KIFU FOREST RESERVE (SITE 9, TOWER 66)

A total of 90 species were recorded in Kifu forest reserve. These belonged to 37 families, of which tree species constituted 27.7% of the total (Appendix P1). Kifu forest reserve has, over the years, been replanted with plantation trees by the National Forestry Authority. The northern side of the existing powerline was replanted with *Araucaria cunninghamii*, *Araucaria haustenii* and *Araucaria agathis* between May 2001 and October 2002. Other species planted on the northern side include *Eucalyptus grandis*, *Eucalyptus odorata* and *Eucalyptus paniculata*. This survey revealed that there was an average stem density of 750 trees ha⁻¹ of *Araucaria cunninghamii* with dbh ranges of 6.3 – 19.7 cm. The southern side was replanted with *Maesopsis eminii* with an average stem density of 400 stems ha⁻¹.




Figure P4: Diameter size classes for the trees found on the southern side of the existing powerline in Kifu forest reserve.

The understorey of the *Maesopsis eminii* plantation is characterised by thick undergrowth and scattered trees of mainly *Funtumia elastica*, *Alchornea cordifolia*, *Antiaris toxicaria* and *Celtis mildbraedii*. The relative diameter size distributions of the trees are given in Figure P4. Stem

densities ranged from 50 stems ha⁻¹ in the 21–30 diameter class to 200 stems ha⁻¹ in the 11–20 diameter class. The understorey is characterised by species such as *Phaulopsis angolana*, *Glycine wightii*, *Marantochloa leucantha*, *Pteris dentata*, *Peddiea fischeri*, *Renealmia congolana*, *Acanthus pubescens*, *Vernonia amygdalina*, *Acalypha bipartita*, *Acalypha ornata*, *Scutia myrtina* and *Rubus apetalus* (Appendix P1).

2.23 NAMYOYA FOREST RESERVE (SITE 10, TOWER 51)

Namyoya forest reserve has, over recent years, been replanted with *Eucalyptus grandis* woodlots and plantations (Plates 12 & 13). However, some remnant indigenous species still exist in a tiny forest patch (about 1 ha.) on the northern side of the existing power line. The southern side also has some indigenous species but many have been cut down to pave way for cultivation or replanting with eucalypts. A total of 43 species belonging to 17 families were recorded in Namyoya forest reserve. The richest family was Asteraceae with 14% of the total number of species (Appendix P1). Some of the indigenous tree species recorded include *Funtumia elastica*, *Alchornea cordifolia*, *Sapium ellipticum* and *Erythrina abyssinica*. The eucalyptus stand on the northern side is in the sapling stage with dbh ranges of 3.8 – 10.6 cm, mean dbh of 6.5 cm and a mean density of 850 stems ha⁻¹.

2.24 SIMILARITIES AMONG MABIRA, KIFU AND NAMYOYA FOREST RESERVES

Based on the presence/absence data of species, the similarity among the three forests investigated, based on the species assemblages, was explored using cluster analysis (Figure P5).




Figure P5: Cluster analysis of the three forest reserves based upon the presence or absence of 315 plant species using a Simple Matching coefficient (SM) cut-off of 0.2.

There was a closer similarity between Namyoya and Kifu forest reserves (0.713) than Mabira. This may be attributed to the relatively high number of species characteristic of disturbed areas that were recorded for Namyoya and Kifu forest reserves. However, it should be noted that the relatively low number of species observed for Kifu and Namyoya forest reserves could also influence the clustering process.

2.25 GENERAL DISCUSSION AND COMPARATIVE OVERVIEW WITH PREVIOUS STUDY

The total number of species (315) recorded for the three reserves compares well with those recorded in the previous survey. However, some of the species recorded previously have not been recorded in the present survey. This is more pronounced with Mabira forest. Species such as *Abrus canescens*, *Basella alba*, and *Strombosia scheffleri* were not recorded in the present study. This may be explained by this spatial rarity since the previous survey recorded them only once on a particular transect. Likewise, there are species that were recorded this time that were not recorded in the previous survey. These include *Psilotricum scleranthum*, *Senecio syringifolius*, *Stictocardia beraviensis* and *Mukia maderaspatana*, which are mainly herbaceous plants. Their being herbaceous and spatially rare, may partly explain why they were not recorded in the previous survey. Some tree species, mainly *Celtis mildbraedii* and *Celtis wightii* were recorded in higher abundances than previously reported in sites 1 and 2, which are dominated by *Broussonetia papyrifera*. Langdale-Brown *et al.* (1964) reported that species of the genus *Celtis* in Mabira forest could represent a pre-climax forest and therefore sites 1 and 2 will, over the years, tend towards a pre-climax stage. There is a considerable variation in species compositions for Namyoya and Kifu forest reserves. This can be attributed to the conversion of the reserves into plantation forests requiring specific silvicultural practices that obviously influence the species compositions.

Given the logging history and disturbance due to encroachers on Mabira forest, the values of basal area recorded are not surprising. Tropical secondary forests, like Mabira, usually undergo rapid accumulation of biomass during the first 15 years or so and then slow down, often reaching values of basal area comparable to those of mature forest before maturity (Brown & Lugo, 1990).

Tree density, particularly for smaller trees, was the most pronounced difference in the vegetation structure of the three cluster groups in Figure P3. Whereas it was expected that sites 1 and 2 would have a higher stem density, given the prolific regeneration of *Broussonetia papyrifera*, clusters M1 and M2 in Figure P3 exhibited a comparable similarity in stem density. This result agrees with a well-known self-thinning process of aging secondary forests in which a declining tree density, mostly caused by mortality rates of smaller trees (< 10 cm dbh), is compensated by the growth of surviving trees (Oliveira Filho *et al.* 1997). Therefore as forest regeneration proceeds, the average tree size increases while tree density declines. However, one cannot rule out the role of harvesting of trees for timber, poles or fuelwood in influencing the richness and diversity of species as observed in some sites sampled in the forest. In general, tropical secondary forests restore species richness first and then diversity, species composition, and finally vegetation structure, particularly tree density, all within a time span between 50 and 150 years (Saldarriaga & Uhl, 1991).

The high basal area recorded on site 3 can be attributed to the presence of a relatively higher number of tree individuals with very high dbh, representing the biggest trees recorded, compared to Site 8. The biggest trees recorded on this site were mainly of the genera *Albizia* and *Alstonia* with dbh up to 170 cm.

2.26 SPECIES OF CONSERVATION CONCERN AND ECONOMIC IMPORTANCE

The Mahogany species namely, *Entandrophrama cylindricum*, *Entandrophragma angolense* and *Khaya anthotheca* are listed as globally threatened and categorized as Vulnerable (IUCN, 2000). These were recorded in Mabira forest in lower diameter classes but only three individuals were observed and they were beyond the planned wayleave extension. They provide first class timber.

Albizia spp are used as second class timber species (Katende *et al.* 2000). Other species commonly used for timber are *Antiaris toxicaria*, *Trilepisium madagascariensis* and *Holoptelea grandis*. Several species are used for medicinal and related purposes including *Citropsis articulata*, *Antiaris toxicaria*, *Albizia grandebracteata*, *Alstonia boonei*, *Bridelia micrantha* and *Croton megalocarpus* (Katende *et al.* 2000). *Dracaena fragrans* is used as an ornamental, hedge and boundary marking. Most of the woody species can be used as poles for local construction, tools, firewood and charcoal burning. Invasive alien plants such as *Broussonetia papyrifera* and *Lantana camara* are a serious threat to plant biodiversity through the formation of very dense populations that affect the population dynamics of the persisting species (Mack *et al.*, 2000). Land use changes such as the replacement of natural ecosystems by agricultural systems, such as Mabira forest encroachment until the early 1990s, alter many ecosystem functions and may promote biological invasions (Hobbs, 2000).

2.27 ALIEN INVASIVE SPECIES: BROUSSONETIA PAPYRIFERA, TITHONIA DIVERSIFOLIA AND LANTANA CAMARA

Sample sites 1 and 2 in Mabira forest were predominantly *Broussonetia papyrifera* whereas the wayleave was dominated by *Lantana camara* and *Tithonia diversifolia* in most areas sampled. The removal of encroachers from Mabira forest in 1992 created large areas of forest land with minimal or no tree cover. This provided suitable conditions for the growth and proliferation of *Broussonetia papyrifera*, a light demander. The continued periodic removal of vegetation in the wayleave has maintained the most prolific and resilient species that are able to withstand the periodic disturbances including *Lantana camara* and *Tithonia diversifolia*. Light is an important plant resource (Blankenship, 2002) that may interact with other plant resources to affect plant performance. Below certain thresholds, however, light limitation alone can prevent seedling survival regardless of other resources (Tilman, 1982). It is therefore probable that the vertical stratification of *Broussonetia papyrifera* may reduce the intensity or duration of light and thus prevent the establishment of other tree species seedlings. Low light has been shown to affect the distribution of other herbaceous species in understorey habitats (Sharma *et al.*, 2005) and this may have important management implications for biological invasions and maintenance of biological diversity.

2.3 WETLAND ECOSYSTEMS – LUBIGI SWAMP

2.31 INTRODUCTION

In Uganda, the rising human population together with increased agricultural production has led to substantial pressure on wetland resources (Mafabi & Taylor, 1993). It is the lowland valley and swamp forest wetlands that have currently come under the strongest pressure. The small valley swamps and seasonal wetlands are closely associated with human activities and it is in these that a heavy toll on resources has occurred (Mafabi & Taylor, 1993). Though these swamps are small in size, individually they are of great significance to the people and to wildlife conservation.

Wetlands are however difficult to classify because they share the characteristics of both permanently wet and generally dry environments. The difficulties are compounded further by the enormous variety of wetland types, and their highly dynamic character, which complicates defining their boundaries with precision (Maltby, 1991). Visser (1960) recognized two types of wetlands in Uganda, the lake edge/Nile shore swamps and the valley swamps. Whereas both may occur in the same region, the lake edge and the Nile shore swamps are more typical of central, northern and eastern Uganda. Carter (1956) also described two kinds of swamps in Uganda,

namely the littoral swamps covering the lakes, and the Nile shores and the shallow valleys, which are characteristic of the country between Lake Albert, Victoria and Kyoga. However, Mafabi and Taylor (1993) classified the Uganda wetlands into three categories, namely: swamps, swamp forest and sites with impeded drainage.

Swamps tend to be dominated for large tracts by a single, vigorous species, which in most cases are sedges (Moss, 1980). *Phragmites*, *Typha* and *Cyperus papyrus* characterise many African swamps (Moss, 1980). Visser (1960) had earlier recognised that 60% of the permanent swamps of Uganda consist of *Cyperus papyrus* which agrees with the more recent view by Moss (1980). Carter (1956) also recognised four succession zones in the swamps of Uganda and observed the presence of members of family Cyperaceae in each zone, with one zone being dominated entirely by *Cyperus papyrus*.

The dominant plant in the Uganda wetlands is *Cyperus papyrus* whose stalks rise, sometimes as a dense forest, some four to five metres above the surface of the rhizomes embedded in a mat of decaying vegetation (Beadle & Lind, 1960). A typical valley wetland adjoining Lake Victoria is fringed with dense forest, an important constituent of which is the wild date palm (*Phoenix reclinata*). The forest does not extend across the lower reaches of the valley, although seedlings are often found there. This, according to Beadle & Lind (1960), may be due to the fluctuations of the water depth as well as to the practice of firing the swamp vegetation in the dry season (see below, and Section 6.4).

Cyperus papyrus is not always the dominant plant in the lower valley wetlands, it is replaced in different parts by the bulrush (*Typha capensis*), *Cladium mariscus* or by *Phragmites mauritanus* (Beadle & Lind, 1960). They further noted that each of these plants seemed to have its own requirements, and the nature of the plant cover could prove to be a valuable indicator of the quality of the mud below with respect to acidity, nutrients, degree of flooding and silting, etc. *Phragmites*, for instance, is common in regions of former volcanic activity where it occurs abundantly in the valleys and around lake edges. *Typha* is found in flooded silted areas and *Cladium mariscus* in less acidic muds.

After *Cyperus papyrus* the commonest wetland plant is *Miscanthus violaceus* (Beadle & Lind, 1960). It often grows around the dry landward edge of the lake-bay wetlands and also forms floating mats. Beadle and Lind (1960) reported that *Miscanthus* seems to favour rather acid conditions and is often found in small lakes surrounded by grasslands on sandy soils and it is sometimes associated with species of *Sphagnum* moss.

Grazing in wetlands especially in the dry season has been a common traditional practice which has survived many generations. This is because these wetlands provide an important source of fresh grazing material in the dry season. Since wetlands have been used to sustain livestock within their proximities in the dry season, they have been subjected to periodic burning in order to encourage the growth of fresh vegetation. Burning has also been applied to these wetlands so as to open up the wetland to access the open water for fishing or to hunt the Sitatunga whose habitat is wetland. Other activities carried out in the wetlands include harvesting of *Cyperus papyrus*, burning, cultivation and grazing (Mafabi & Taylor, 1993).

2.32 METHODS

Transects of up to 200 m were used, following habitat types that characterized each site as much as possible (Figure P6). Line transects and circular plots were used to collect the species richness data. Sampling points were located at 10 m intervals. Using the sampling point, two circular plots

of 1 m radius were located equidistant from the sampling point and from each other at a distance of 2 m. This is a modification of the point frequency method (Okland, 1990) and the method used by Wettstein & Schmid (1999). All species found in these plots were recorded and those that could not be identified in the field were collected for identification at the Makerere University herbarium.

To place sites into meaningful groups, cluster analysis was used to produce a dendrogram containing all the six sites using the agglomerative clustering technique provided in the Multivariate Statistical Package (MVSP) of Kovach (1999). A minimum Simple Matching Coefficient (SM) of 0.2 (dotted line in Figure P6) was used for defining clusters.

2.33 RESULTS

Floristics

There were 124 species, belonging to 90 genera and 36 plant families. The most dominant plant family was Fabaceae with 20 species followed by Cyperaceae, Poaceae and Asteraceae with 17, 12 and 12 species respectively (Appendix P2). The most abundant genus was *Cyperus* with six species followed by *Scleria* with five. The tree species constituted 13% (16 species) of the total species while the herbs represented 63% (78 species). The commonest herbaceous species recorded were *Pycnus nitidus*, *Panicum maximum* and *Cyperus denudatus* whilst the commonest tree species was *Phoenix reclinata*, which was recorded at five of the six sites surveyed. Among the rare but typical wetland species were *Enhydra fluctuans*, *Eulophia horsfallii*, *Geniosporum rotundifolium*, *Lygodium microphyllum*, *Lysimachia ruhmeriana*, *Mikania cordata*, *Nephrolepis biserrata*, *Polygonum strigosum*, *Siegesbeckia abyssinica*, *Stephania abyssinica*, *Gomphocarpus fruticosus*, *Ficus verruculosa* and *Tabernaemontana odoratissima*. Each of these was found in only one of the six sites.

2.34 SIMILARITY AMONG SITES

Cluster analysis (Kovach, 1999; Okland, 1990) was used to determine the degree of similarity among sites based on the presence or absence of species. Two distinct community types were easily identifiable as shown in Figure P6. Cluster Sw 1 consists of sites with an extensive seasonally flooded area, subjected to grazing. The characteristic species of these sites include open grassland species such as *Loudetia kagerensis*, *Scleria melanomphala*, *Sporobolus pyramidalis*, *Eragrostis racemosa*, and *Indigofera* sp. Other species that represent swamp forest and thickets include *Maesa lanceolata*, *Rhus* sp., *Alchornea cordifolia*, *Leersia hexandra*, *Macaranga* sp., and *Phoenix reclinata*. Areas with semi-permanent and permanent water are characterised by *Loudetia phragmatoides*, *Cyperus papyrus*, *Miscanthus violaceus* and *Kotschyia africana* as the dominant species.




Figure P6: Cluster analysis of the six sample sites in Lubigi swamp based upon the presence or absence of 124 plant species. Site 1 (Kazinga 1), Site 2 (Kazinga 2), Site 3 (Lubanyi 1), Site 4 (Lubanyi 2), Site 5 (Nakawudde) and Site 6 (Nganda-Nansana).

Cluster Sw 2 is characterised by thick *Cyperus papyrus* stand that are less species rich compared to sites in cluster Sw 1. The swamp fringes are characterised by *Acanthus pubescens*, *Phoenix reclinata*, *Aframomum angustifolium*, *Urena lobata*, *Teramnus* sp., *Leersia hexandra*, *Bridelia micrantha*, *Mimosa pigra*, *Alchornea cordifolia* and *Bothriocline bagshwei*. These then give way to the permanent deeper water tolerant species such as *Hyptis lanceolata*, *Ficus verruculosa*, *Cyperus papyrus*, *Typha domengensis*, *Eulophia* sp. and *Cyclosorus striatus*.

3.0 INVERTEBRATES: BUTTERFLIES AND DRAGONFLIES

3.1 INTRODUCTION

Approximately 200 species of butterflies and 60 species of dragonflies are known to occur in Mabira forest (Davenport *et al*, 1996; JJ Kisakye, *pers comm.*). The butterfly and dragonfly fauna of Kifu forest, Namyoya forest and Lubigi swamp have not been previously documented.

Little research has been conducted on the impact of transmission lines on terrestrial insects and other arthropods. The impact of clear-cutting the forest along transmission lines is limited since most invertebrate species adapt quickly and migrate to either the clear-cut, mature forest/edge, or interior forest areas.

This study was mainly carried out to document the butterfly and dragonfly species from the different sites covered by this project. The study was intended to

- 1 document species along the proposed transmission line
- 2 provide a description of the anticipated environmental effects
- 3 identify potential impacts on the invertebrate diversity by construction of the transmission line.

3.2 METHODS

The butterfly and dragonfly fauna of Mabira was sampled through the systematic use of sweep nets and baited traps (in the case of butterflies) for a total of 8 man-days. Fermenting banana was used as bait for the traps. For the other sites, only intensive use of sweep nets was employed. Sweep netting was done both along an established 0.5km transect line as well as random sweeping within the entire areas around each sampling site. This approach also known as Rapid Biodiversity Assessment, involved combing through the entire area, and catching every species encountered. A number of standard field guides were used, as well as the extensive collections at the Zoology Museum, Makerere University, for identifying any butterfly specimens that were collected. Preliminary identifications of common and familiar butterfly species were done in the field. Mr. Joseph Kisakye (Department of Zoology) helped with the identification of the dragonfly specimens, using a number of keys.

The sampling locations that were used are detailed in Appendices G1 and G2.

3.3 RESULTS: DRAGONFLIES

20 species (Appendix A1) were recorded, including three belonging to the suborder Zygoptera and 17 species belonging to Anisoptera. Together, these account for only 8.6% of Uganda's species. For the purposes of this report, we refer to all members of the Order Odonata as 'dragonflies'.

A total of 17 species were recorded for Mabira forest, accounting for 29% of the known species for this forest. A few taxonomically difficult species were only identified to generic level, meaning that the total diversity of species recorded is a little higher than indicated here.

Two particular species are worth highlighting: *Orthetrum macrostigma* was only previously known to occur in Bwindi forest (JJ Kisakye, *pers.comm.*) and the Uganda endemic species, *Chlorocypha trifaria*; both recorded in Mabira forest in site 3. This site also registered the highest species richness, with 8 out of the 17 recorded. In four of the sites dragonflies were recorded.

Namyoya and Kifu forests had very low dragonfly species richness with only three and four species respectively. This may be attributed to the status of the forests that are quite degraded and with few wetlands compared to Mabira.

From Lubigi swamp on the other hand, a total of 9 species were recorded, with sites 1 and 3 registering 5 species each. Site 6 had the highest total number of species with 6, whilst site 4 registered only 2 species.

Species abundance

Based on relative commonness of individuals seen, Table A1 presents a scheme that has been used to categorize the species abundance, which varied both between and within sites.

Table A1. Dragonfly species abundance categorization

Number	Number of individuals observed	Range category
1	1	Rare
2	2-3	Relatively rare
3	4-5	Relatively abundant
4	6-9	Abundant
5	>10	Very abundant

For Mabira forest, 6 out of the 17 species fall into category 1, accounting for 35.3% of total species recorded. Eight species belonged to category 2 (47.1%) and only three species belonged to category 3. None of the species could be said to be very common in Mabira forest, although this picture may not hold for other areas.

3.4 RESULTS: BUTTERFLIES

The total number of species recorded were 165 (Mabira), 65 (Kifu FR), 48 (Namyoya FR) and 56 (Lubigi) respectively (Appendix A2). Table A2 summarises the numbers of species recorded in the different families for the different areas. For Mabira this number is less than that recorded during the 1990s biodiversity surveys that registered 199 species in total. However, an unpublished report puts the number higher at 218 species for this forest.

Table A2. Summary of species recorded by this study

Family	Mabira				
	This study	Previous records			
Nymphalidae	102	128	41	32	28
Pieridae	23	24	5	2	7
Lycaenidae	17	25	11	4	8
Papilionidae	10	9	4	2	2
Hesperiidae	15	13	6	8	11
Total	167	199	67	48	56

For Mabira Forest Reserve, there was a slight increase in number of species recorded from three families: Pieridae, Lycaenidae and Papilionidae as compared to those reported in Davenport *et al.* (1996). The other families however registered declines in numbers of species, with Nymphalidae having 26 species less.

3.5 DISCUSSION

Five restricted range species (Davenport *et al*, 1996) were recorded during this study, namely *Neptis trigonophora* (179) F and *Acraea rogersi* (179) F that were only recorded in Mabira. The others were recorded in 2 or 3 forests: *Acraea aganice* (174) f, *Celaenorrhinus intermixtus* (169) F and *Celaenorrhinus bettoni* (169) f.

Appendix A3 compares the species of butterflies previously recorded with my surveys. From the appendix, it can be seen that 34 species of butterflies that were previously recorded from Mabira forest have not been recorded by this study. On the other hand, 50 species not previously known for Mabira have been added to the existing species list. I attribute the additional number of species to the ability of more open and widespread species to colonize areas of disturbances within the forest

Table A3. Summary of butterfly species with their respective ecological types according to Davenport *et al* (1996)

Ecotype (see text)	Sites			
	Mabira	Kifu	Namyoya	Lubigi
F	92	29	2	5
f	26	18	15	11
O	6	2	4	8
W	27	14	22	18
S	0	0	0	5
M	16	4	5	9

Every species recorded by this study was assigned to one of the ecological categories: F-forest dependent species, f- forest edge/woodland species, S- swamp/wetland species, O- open habitat species, W- widespread, M- migratory species (Table A3). The proportion of the F and f species in a sample is used as an indication of the ecological state of the habitat. For Mabira forest, 71.6% of total species fell in these categories, with 55.8% F- species. Only 29.7% of the butterfly fauna belonged to the O, S and W ecotypes. This picture however shifted dramatically in the much more degraded Namyoya forest where only 4.2% of the butterfly fauna are forest dependent. Over 50% of the species in Namyoya are open habitat dwellers and/or widespread.

Table A4. Butterfly species abundance data.

Number	Number of individuals	Rank category
1	1-3	Rare
2	4-10	Relatively rare
3	10-25	Relatively abundant
4	25- 39	Abundant
5	≥40	Very abundant

For all four sites, only 5 species can be described as rare based on the ranking in Table A4. These were *Papilio nobilis*, *Mimeresia sp*, *Charaxes zelica*, *Neptis ochracea* and *Iolais pasasilanus*. Most of the other species fell in category 3 or 4 and only a few species were recorded for category 5. In Mabira, most of the species were relatively rare especially within the deeper and denser forest areas at sites 5 - 8. For Kifu and Namyoya forests, the numbers were much lower.

4.0 AMPHIBIANS AND REPTILES

4.1 INTRODUCTION

Both Amphibians and Reptiles are classes of vertebrates, the forest comprising of frogs, toads, caecilians, newts and salamanders. In their evolution amphibians are said to have evolved from fish 350 million years ago (Young, 1981). Reptiles Include turtles and tortoises, lizards, chameleons, skinks including limb-less skinks, geckoes, crocodiles, monitors and all types of serpents commonly known as snakes (Foster, 1994; Goin *et al*, 1962). Reptiles are said to have evolved from amphibians some 270 million years ago.

Ecologically, amphibians are important in many ways; they are mostly predators, acting as primary and secondary carnivores. Their prey consists mostly of insects, some of which are pests of crops or disease vectors. Amphibians are therefore important ecological components of both wetlands and dryland. Among vertebrates they are distinctive in many ways. A thin, moist, highly permeable skin; jellied, unshelled eggs; possession of aquatic and terrestrial life histories; restricted home range; and limited dispersal abilities of many species make amphibian effective biomonitors. For biological assessments, they are especially promising because of their capability of linking wetlands with surrounding landscapes (upland habitats) (U.S. EPA. 2002). They are also interlinked in food chains, often acting as food for other vertebrates, such as pigs, birds, snakes and sometimes man. Because of their ectothermic physiology, the life history and ecology of amphibians often differ markedly from that of birds or mammals (McCollough *et al*, 1992).

Reptiles are also ecologically important. They feed on a number of animals and this predation involves reptiles in ecosystem food webs.

The area where the proposed Bujagali Hydropower line is to pass or to be erected is also inhabited by the herpetofauna. This study was designed to find out the status of the amphibian and reptilian species and whether the proposed activities would have significant effects on their habitats and the species themselves.

4.2 OBJECTIVES

The overall objective was to study amphibians and reptiles in the areas where the proposed power line for Bujagali Hydropower project will pass. It also involved evaluating the potential impacts. Specifically, the work involved:

- Generating species lists for amphibians and reptiles in and around the proposed area where the power line will pass.
- Identifying the species of conservation concern.
- Determining the impact that the construction of the power line would have on them.

Impacts on the habitats where amphibians and reptiles are found are reflected in changes in numbers and species diversity in a short time. These are some of the factors that have led to amphibians being recognized today as good indicators of habitat change. The geographical ranges of amphibians are smaller than those of other vertebrates (Bibby *et al*, 1992), which suggest that amphibians are more likely to be affected by habitat changes than other vertebrates.

Most reptiles on the other hand are highly mobile and are diversified in habitats. They can be encountered in aquatic habitats, shorelines, rock outcrops, trees and bushes and on any slopes of

the terrain. Those more specialized in habitat use such as crocodiles, monitor lizards and water snakes are good indicators for monitoring changes in a habitat due to human activity.

Study Area

The surveys for the amphibians and reptiles were conducted in the locations detailed in Appendices G1 & G2. Additional locations that were particularly important for amphibians were also noted even when they lay outside the eight sample sites in Mabira, but were along the general transmission line area.

4.3 METHODS

Several methods are available for surveying amphibians (reviewed in Heyer *et al*, 1994; Fellers and Freil, 1995; Halliday, 1996; Olson, *et al*, 1997). These include visual encounters, egg surveys, and call surveys, terrestrial cover boards, dip nets, seines, aquatic funnel traps, and terrestrial pitfall traps. The method is dictated by the habitat type. Because of the time limitations, three sampling methods were employed during study, namely opportunistic observations, visual encounter surveys and acoustic Surveys.

Opportunistic Observations/Searches

Opportunistic searches were used to maximize the number of species encountered in the study area. This method involved recording any amphibian or reptilian species encountered anywhere and at any time in the study area, or brought in or reported by local people.

Visual Encounter Surveys

The visual encounter survey (VES) method is commonly used to determine the species richness of an area, to compile a species list and to estimate relative abundances of species within an assemblage. It was used to determine the species richness of the study areas. This involved walking through the sampling areas or habitat for a prescribed time period systematically searching for amphibians.

The VES was done along the transects established by the research team for the purpose of studying other taxa, and also along the streams and ponds, sampling all amphibians and reptiles that were visible. This focused on surface-dwelling amphibians and reptiles.

Local Consultations and Literature Review

Local people can be a valuable source of information. Some are constantly in touch with their environment, encountering amphibians and reptiles of different kinds as they carry out their activities. Talking to the local people yielded one species record of family Testudinidae. The record was confirmed by reviewing literature of studies carried by other researchers.

Data Analysis

The reptiles and amphibians were identified using standard reference books available namely; Schiotz (1972), Schiotz (1972b), De Witte (1937), Drewes (1984), Drewes and Vindum (1994), Loveridge, (1957), Welch (1982), Stewart (1967), and Wager (1965). Kigoolo (1994) and Behangana (1995) were also useful for comparison of species' distribution in Uganda.

Unlike other taxa such as birds or mammals, no standard species list exists, although the Herps Working Group of Nature Uganda is developing one. However, species categorization using the IUCN Red Data Book categories for some species is available and has been used.

4.4 RESULTS

A total of 19 amphibian species and 6 reptilian species were recorded present during the survey as shown in Tables H1 – H4.

4.41 AMPHIBIANS

Eight (8) amphibian species were recorded in Mabira Forest reserve; while sixteen (16) were recorded in Lubiji Swamp. Most of the species recorded in Lubiji swamp are wetland specialists. Only two species (*Arthroleptis adolfifriederici* and *Leptopelis christy*) recorded present in Mabira Forest reserve are purely forest specialists.

The commonest species were members of family Hyperoliidae (genera *Afraxalus* and *Hyperolius*) followed by family Ranidae (genera *Ptychadena*, *Phrynobatrachus*, *Afrana* and *Hoplobatrachus*). Family Pipidae (genus *Xenopus*) was also common especially in Lubiji (Kazinga 1, sampling site). More specifically, members of family Hyperoliidae found to be most common include *Afraxalus quadrivitatus*, *Hyperolius cinnamomeoventris*, *Hyperolius viridiflavus bayoni* and *Hyperolius viridiflavus viridiflavus*. Those of family Ranidae included *Ptychadena mascareniensis*, *Phrynobatrachus natalensis*, *Afrana angolensis* and *Hoplobatrachus occipitalis* in that order.

The more individuals of a given species you encounter are indicative of the abundance of that particular species. *Afraxalus quadrivitatus*, *Hyperolius cinnamomeoventris*, *Hyperolius viridiflavus bayoni* and *Hyperolius viridiflavus viridiflavus* are therefore the most abundant amphibian species. These are followed by *Ptychadena mascareniensis*, *Phrynobatrachus natalensis*, *Afrana angolensis* and *Hoplobatrachus occipitalis*.

The said common species of family Hyperoliidae are generally associated with permanent water sources and tend to select habitats with water all year round. These were mainly recorded in Lubiji Swamp sites. Several of these species, however, have the ability to resist temporary and regular drying up of their habitats (Dudley, 1978). Members of the genus *Hyperolius*, commonly known as Reed frogs are most active from dusk to dawn when they are highly active but they are hardly noticed during the day time. They are commonly found perched on swamp wetland vegetation such as water reeds and papyrus.

Tables H1 & H2 present the lists of amphibians recorded in Mabira CFR and Lubiji wetland sites.

Members of genera *Xenopus*, *Afrana* and *Hoplobatrachus* are also associated with permanent water sources. They are commonly found near water, more so for the bullfrog which only gets out of water to feed. *Afrana angolensis* is a riverine species found mainly along rivers and this was encountered along rivers in Mabira Forest Reserve. *Xenopus* is more aquatic than the rest and is found in water most of the time. *Xenopus* and the bull frog were mainly recorded in Lubiji swamp, on swamp edges and along transects in the swamps.

One member of family Arthroleptidae, *Arthroleptis adolfifriederici* was recorded for the first time in Mabira Forest Reserve. No threatened species were recorded.

Table H1: Amphibian encountered in Mabira Forest Reserve

Family	Species	Common name	Total number of Individuals
Hyperoliidae	<i>Leptopelis christy</i>		10
Pipidae	<i>Xenopus laevis</i>	African Clawed Frog, Common Platanna	2
Ranidae	<i>Phrynobatrachus acridoides</i>	Eastern puddle frog	1
	<i>Phrynobatrachus natalensis</i>	Natal Dwarf Puddle Frog	9
	<i>Ptychadena anchietae</i>	Anchieta's ridged frog, Plain Grass Frog	1
	<i>Ptychadena mascareniensis</i>	Mascarene ridged Frog	15
	<i>Afrana angolensis</i>	Angola river frog	11
Arthroleptidae	<i>Arthroleptis adolfifriederici</i>	?	1
Totals	8 species		

Table 2: Amphibian encountered in Lubiji Swamp

Family	Species	Common name	Total number of Individuals
Bufonidae	<i>Bufo reguralis</i>	African Common Toad	4
	<i>Bufo maculatus</i>		7
	<i>Afrixalus quadrivitatus</i>	Four-lined Leaf-folding frog	30
	<i>Hyperolius kivuensis</i>	Kivu Reed Frog	10
	<i>Hyperolius cinnamomeiventris</i>	Cinnamon-bellied reed frog	26
	<i>Hyperolius viridiflavus bayoni</i>		29
	<i>Hyperolius vidiflavus viridiflavus</i>		32
	<i>Kassina senegalensis</i>	Senegal Kassina	12
	<i>Leptopelis christy</i>		1
Pipidae	<i>Xenopus laevis</i>	African Clawed Frog Common Platanna	10
	<i>Phrynobatrachus natalensis</i>	Natal Dwarf Puddle Frog	4
	<i>Phrynobatrachus graueri</i>		3
	<i>Ptychadena anchietae</i>	Anchieta's ridged frog, Plain Grass Frog	2
	<i>Ptychadena mascareniensis</i>	Mascarene ridged Frog	8
	<i>Ptychadena oxyrhynchus</i>	Sharp-nosed ridged Frog	1
	<i>Hoplobatrachus occipitalis</i>	Crowned bullfrog	11
Totals	16 species		

4.42 REPTILES

Five reptile species were recorded in Mabira and Kifu Forest Reserve and five in Lubiji swamp. Of special interest are the Nile Monitor (recorded in Kifu Forest) and the African Rock Python recorded in Kifu Forest and Lubiji swamp (Site 1). The two species are of importance to the people in the area.

Reptiles, like amphibians are cold-blooded vertebrates. They utilize the sun's energy to raise their body temperatures in order to be more active. Therefore, the best sampling time for reptiles are the early hours of the day when they come out of hiding to bask. The commonest reptilian species was Jackson's Lizard, *Lasutus jacksoni* with 16 individuals encountered followed by the Blue-headed Agama *Agama atricolis* for which 5 individuals were encountered. Both these species were encountered within Mabira forest reserve and on culverts along access road in Mabira forest.

One species of family Testudinidae, the soft-shelled turtle, was reported by the local community at Ganda sampling site (Site 6) in Lubiji swamp.

The species recorded in the different study areas are listed in Table H3 for Mabira and Kifu CFRs & Table H4 for Lubigi swamp.

Table H3: Reptilian fauna recorded from Mabira & Kifu Forests

Order Squamata	Species	Common Name	Total number of individuals
Mabira Forest Reserve			
Family Lacertidae	<i>Lasutus jacksoni</i>	Jackson's Lizard	14
Family Agamidae	<i>Agama atricolis</i>	Blue-headed Agama	4
Suborder Serpentes			
Family Colubridae	<i>Rhamnophis aethiopissa</i>	Large-eyed Tree Snake	1
Kifu Forest Reserve			
Order Squamata			
Family Varanidae	<i>Varanus niloticus</i>	The Nile Monitor	1
Suborder Serpentes			
Family Pythonidae	<i>Python sebae</i>	Rock Python	1
Total	5 Species		

Table H4: Reptilian fauna recorded from Lubiji Swamp

Order Squamata	Species	Common Name	Total number of individuals
Family Lacertidae	<i>Lasutus jacksoni</i>	Jackson's Lizard	2
Family Agamidae	<i>Agama atricolis</i>	Blue-headed Agama	1
Suborder Serpentes			
Family Pythonidae	<i>Python sebae</i>	Rock Python	1
	<i>Naja melanoleuca</i>	Water Cobra	1
Order: Testudinidae			
Family Trionychidae	<i>Trionyx triunguis</i>	Soft-shelled Turtle	0
Total	5 Species		

4.43 SPECIES OF CONSERVATION CONCERN

Certain species in wetlands merit special attention due to their scarcity, restricted ranges or unique habitat requirements. In Uganda, among the amphibians, only frogs and toads have been recorded. However, like in many other developing countries of the world, amphibians of Uganda are less known than reptiles, birds and mammals. The conservation status of amphibians in Uganda is generally unknown because of data deficiency.

As for reptiles, no species of conservation concern were encountered or recorded in the areas to be traversed by the proposed where we conducted our surveys. .

4.5 DISCUSSION

In Uganda, among the amphibians, only frogs and toads have been recorded. However, as in many other developing countries, amphibians are less known than reptiles, birds and mammals. The conservation status of amphibians in Uganda is generally unknown because of data deficiency. No reptiles of conservation concern under the IUCN red list categories was encountered.

The amphibian and reptile species encountered during the limited surveys for this report, are widely distributed in Uganda. The majority, especially those in the wetland are quite abundant. The population to be affected by the construction of the power line is therefore small and will not affect the overall survival of the species. No species of conservation concern were recorded in the areas where the power line will pass.

5 BIRDS

5.1 FOREST BIRDS: INTRODUCTION

The birds of Mabira Forest are well-known, from the work of Carswell (1986), Davenport *et al* (1996), Rossouw and Sacchi (1998) and Naidoo (2003).

5.2 METHODS

Most data were collected along transect lines, as described in section 1.2 and Appendix G1. Birds were recorded opportunistically at any time, but most data derive from two standard methods. The first, *Timed Species Counts (TSCs)*, are described in the literature (Freeman *et al*, 2003) as well as in the previous report (AESNP, 2001) and our 2006 Bujagali report (Pomeroy, *et al* 2006). Essentially, all birds seen or heard (and the latter predominate in forests) are recorded in the order in which they are detected, and scored out of six (for the first ten minutes), with the score decreasing step by step to 1 for the final ten minutes of a one-hour count. In this study, time was limited, counts were relatively few, and so the results are simply given as averages. In Mabira, two counts were done on each of the transect in 2001, whilst in 2006 only one TSC was done at each transect line. *Mist nets* catch birds because they are so fine that, when set against a dark background, as in a forest, they are almost invisible. They are set in lines along the transects, and checked periodically. Birds caught in the nets are identified and released. Since the nets are set vertically, reaching a height of about 2 metres, they predominantly catch the low-flying birds of the forest understorey – which is where many species mainly live. Altogether, the trapping effect at each transect totalled 1440 metre-net-hours.

As with the Bujagali studies, we categorised birds in various ways, as indicated in Table B1. Of particular importance are the forest specialists, FF and to some extent the generalists, F; the former cannot survive without forest. The ‘f’ species are those that only occasionally visit forest, and typically only forest edge. Collectively we refer to these three categories as ‘tree species’.

Table B1. Bird descriptors. No globally-threatened species were observed. The species’ preferred habitats and migratory status are also indicated in Table B4 and Appendix B1.

Threat categories	G-EN	globally endangered
	G-VU	globally vulnerable
	R-VU	regionally vulnerable
	R-NT	regionally near threatened
	R-RR	species of regional responsibility
Habitat	E	papyrus endemic
	e	Papyrus near-endemic
	W	waterbird
	w	bird often found ear water
	FF	forest specialist
	F	forest generalist
	f	Forest visitors
Migrants	G	species characteristic of grassland
	P	palearctic migrants
	A	afrotropical migrants, migrating within Africa

Global threat categories (G-) are from IUCN’s web site and regional categories (R-) are from Bennun and Njoroge (1969).

5.3 RESULTS AND DISCUSSION

The data from both 2001 and 2006 for all three forests are summarised in Tables B2 - B4, and a more detailed set of the data from 2006 is given in Appendix B1 and B2. The forest is within easy reach of Kampala, and is popular with bird watchers. Birds also featured in the 2001 EIA Report which included information on the smaller forests too. As mentioned in the Introduction (section 1.1.1), the surveys in 2006 were intended, so far as possible, to replicate those of 2001, but some differences were inevitable. To start with, the 2006 studies were in July, whilst those of 2001 were in February, normally a drier month with fewer leaves on the trees and hence better visibility of birds in the trees. A few Palearctic migrants were recorded in 2001, but in July 2006 they will have been breeding in Europe or Asia. During 2001, the main method was Timed Species Counts, complemented by Point Counts. TSCs were also the main method in 2006 but in this case were supplemented by mist-netting, which is an excellent method for hard-to-see understorey birds. Finally, the field work in 2001 was by Isaiah Owionji and in 2006 by Eric Sande. Both, however, are experienced observers of forest birds.

Table B2. Summary of numbers of ‘tree birds’ and other species recorded in Mabira Forest in 2001 and 2006. There were 16 counts in 2001, and 8 in 2006.

Category	2001	2006	Davenport <i>et al</i> (1996)
FF	57	31	75
F	48	31	63
F	38	12	56
Others	37	7	95
Total	180	81	287

Table B3. Summary of ‘tree birds’ and other species recorded in Kifu Forest in 2001 and 2006. In 2001, two TSC counts were made, with three in 2006.

Category	2001	2006
FF	10	7
F	22	17
F	3	11
Others	3	9
Total	38	41

Table B4. Summary of ‘tree birds’ and other species recorded in Namyoya Forest in 2001 and 2006. Two counts were made in 2001, but only one in 2006.

Category	2001	2006
FF	10	1
F	22	7
F	3	14
Others	3	12
Total	38	34

There were substantially fewer forest birds recorded in 2006 than 2001. This is partly explicable in terms of the fewer counts in 2006, as well as the different months of field work, and some

differences in methods resulting from the seasonal change. However, a drop in forest specialist species from 57 to 31, and from 105 to 62 for the combined forest species (FF, F), does suggest that other factors may be involved. We have data from another forest in the area, Ziika, which is much smaller – in fact it is only 14 hectares. Number of species in the two categories combined also dropped, although less steeply, from 23 in 2001 to 17 in 2006 (NBDB, unpublished data). These slightly alarming results should cause us to investigate this further. In neither Mabira nor Ziika does there seem to have been any major change in the habitat during this period. In Mabira in 2001, 72% of the species from the forest as a whole by Davenport *et al* (1996) were noted, compared to only 45% in 2006.

But despite the drop in species numbers, we recorded almost as many species of conservation concern in 2006 as in 2001. For the earlier year, Table 5 listed 32 such species that are known from Mabira, of which only 8 were recorded then. Seven species of conservation concern were recorded in 2006, including two that are globally-listed (Table B5).

Table B5. Globally and regionally threatened species recorded in Mabira Forest in 2006.

Atlas No.	Species	Forest category	Red Data listing
86	BROWN SNAKE-EAGLE <i>Circaetus cinereus</i>		R-NT
124	CROWNED EAGLE <i>Stephanoaetus coronatus</i>	FF	R-VU
156	NAHAN'S FRANCOLIN <i>Francolinus nahani</i>	FF	G-EN, R-VU
290	GREY PARROT <i>Psittacus erithacus</i>	FF	R-NT
498	WHITE-HEADED SAW-WING <i>Psalidoprocne albiceps</i>	f	R-RR
551	TORO OLIVE GREENBUL <i>Phyllastrephus hypochloris</i>	FF	R-VU/RR
559	GREEN-TAILED BRISTLEBILL <i>Bleda eximia</i>	FF	G-VU

The two TSCs in the present wayleave area of Mabira forest yielded 39 species (Appendix B2), including a few forest birds flying over.

Neither of the two smaller Forest Reserves (Kifu and Namyoya) contain any significant areas of natural forest (see Sections 2.122 & 2.123); hence the relatively small numbers of forest birds (see Table B2 and B3, and Appendix B2). Although there were again differences in the numbers of counts at each site, the total numbers of species recorded were similar. The drop in numbers of FF species recorded, especially at Kifu, is therefore likely to reflect the continuing loss of forest trees there.

The degradation of Kifu forest is well-illustrated by the decline in forest specialist (FF) species. Dranzoa (1990) recorded 41, compared to ten in 2001 and a mere seven in 2006.

5.4 SWAMP BIRDS

Unlike the forest sites, swamp birds were not sampled in 2001. The six study sites for birds were the same as for other taxa. A single one-hour TSC (section 3.2) was made at each of the six points listed in Appendix G2. The results, given in detail in Appendix B3 and summarised in Table B6, show that these areas contain a number of species of interest, including three species which are largely confined to papyrus swamps – the so-called ‘papyrus endemics’ (Britton 1978, Byaruhanga *et al*, 2001, Carswell *et al*, 2005, Maclean *et al*, 2006). The best-known of these, the Papyrus Gonolek, is a striking bird of brilliant red and black with a conspicuous yellow crown. Other papyrus birds are less remarkable in their appearance, but because of their restricted distributions, they are considered to be of conservation concern. Two of the four other papyrus endemics, Carruther's Cisticola and White-winged Warbler, were also common. The remaining

two – the Papyrus Canary and Papyrus Yellow Warbler - were not recorded, although the first of these might be found with more thorough surveys. The Greater Swamp Warbler, although not entirely confined to papyrus swamps, can be considered a near-endemic; it was seen at two sites.

The data in Table B6 are divided into two categories: those dominated by papyrus and more open sites with large areas that are flooded only seasonally (clusters Sw2 and Sw1 respectively in Figure P6).

The full species list is given in Appendix B3. The majority of the 94 species recorded are not strictly waterbirds, their presence being due either to their being generalists, such as the Common Bulbul, or to the fact that there were many trees in and near to the swamp. The latter explains the quite large number of ‘tree birds’ (F and f in Table B6). No species of global conservation concern was recorded.

The only species to occur at all six sites was the Grey-capped Warbler, which also had the highest mean score (Appendix B3); next commonest were the Papyrus Gonolek and Winding Cisticola. The average numbers of species per count were high, averaging 32.3 in the more open sites (W3, 4 and 5) and 34.3 at the sites with more vegetation. We used a Jack-knife estimate for total species richness, and obtained a figure of 131, which is also quite high and would probably increase with further sampling. So it is evident that the swamps, despite their closeness to Kampala, and levels of human disturbance, still have notable numbers of birds, of great variety and in good numbers. The ‘papyrus endemics’ are known to be fairly tolerant of moderate levels of disturbance (Maclean *et al*, 2006). There were few species of conservation concern, but the high diversity is in itself a measure of the importance of these swamps.

The more specialised species – and thus those which are of some conservation concern – were quite numerous in the papyrus-dominated sites, and still common in the more open sites (Table B6, last line).

Table B6. Summary of bird data from the six swamp sites (as listed in Appendix G2). The categories of bird types are listed in Table B4.

		Numbers of species	
		Papyrus sites ^a	Open sites ^b
Red Data species	R-VU	1	0
	R-NT	4	0
	R-RR	6	6
Papyrus endemics ^c	E	3	3
	e	1	1
Water birds	W	18	9
	w	16	15
Tree birds	FF + F	6	8
	f	16	24
Migrant status	P	0	4
	A	4	4
Total species of conservation concern ^d		36	27

Notes: a Sites 1, 2 and 6

b Sites 3, 4 and 5

c See Section 4: P indicates papyrus endemics, and p are other species for which papyrus is a major habitat

d Categories E, W, R-VU, R-NT and R-RR

6.0 MAMMALS

6.1 INTRODUCTION

The results presented in this report have been collected from surveys conducted for the same purpose as those we presented in the AESNP (2001) report. In the present study I aimed at conducting mammal surveys to answer similar questions: -

- (i) Documentation of the diversity of mammals in the area along the transmission line
- (ii) Reviewing the relative abundance of species documented through trapping
- (iii) Reviewing the conservation status of the mammal species recorded and
- (iv) Assessing potential and/or real threats to the mammal species recorded along the transmission line.

In addition however the present study.

- (v) Assessed any changes in mammalian diversity and abundance between the two studies

6.2 METHODS

These followed closely the approaches used in AESNP (2001) to enable acquisition of a data set that would enable comparisons to be made between the two studies. We still maintain two groups of mammals (large and small), which require different strategies for their inventory.

The presence of larger mammals (Primates, Carnivores and Ungulates) was recorded through:

- (i) Direct observation for the diurnal species
- (ii) Indirect cues for their presence (such as spoor/paw/foot prints) and fecal pellets
- (iii) Recovery of skeletal material of species, and
- (iv) Interviews with local people.

The smaller mammals (rodents, insectivores and bats) were surveyed using traps or mist nets set along transect established at the same locations used for the surveys in the 2001 study. The nomenclature of these sampling locations is presented in Appendix G1.

For the terrestrial small mammals (rodents and shrews), traps were used to capture them for subsequent identification. Baited traps were used along each of the transects to sample the diversity and abundance of species present in the different locations.

Unlike the study in 2001, this time round, the bats were surveyed at all the transect points and at an additional two locations along the way leave. Well as the results from the netting for bats contribute to the total species list, they would also contribute to the picture of spatial occurrence of the different species if it were not for the selective sampling of the bat fauna by mist nets.

6.3 RESULTS

6.31 MABIRA FOREST

Table M1 summarizes the trap and net success along the various sample transects in the 8 sites. These values represent the quotient of number of individuals captured and the effort invested to capture them.

Table M1 Trap and netting success recorded at the different sampling locations

Transect	Trap success	Net success
Site 1 (at Pylon 179)	4.8	0
Site 2 (at Pylon 174)	2.4	22.2
Site 3(between Pylon 170 & 169)	22.8	37.0
Site 4 (at Pylon 164)	15.8	0
Site 5 (at Pylon 158)	22.9	16.7
Site 6 (at Pylon 154)	24.4	22.2
Site 7 (at Pylon 149)	22.2	11.1
Site 8 (at Pylon 144)	10.4	27.8

Trap success is a fair measure of relative abundance of the species for the terrestrial small mammals (rodents and shrews) although in some situations it may have species that are ‘trap happy’ overrepresented in the results, than compared to those that do not ordinarily easily go into traps. Nevertheless, trap success figures can still give a good indication of relative abundance of the small mammals while for bats these figures may be misleading because certain groups of bats (Megachiroptera or the fruit bats), are easier to capture in nets than the Microchiroptera.

For the terrestrial small mammals therefore sites 1 & 2 returned very low values emphasizing the low abundance of individuals in these two areas of the forest which are primarily composed of *B. papyrifera* with a very sparse undergrowth. The rest of the transects on the other hand are located in more or less intact natural forest which provides more diverse niches and presumably resources for the occurrence of higher numbers of individuals of the different species.

A total of 35 species of mammals have been recorded altogether, which represents 3 species less than those recorded in 2001. The difference however lies in the species composition of species recorded.

The present study represents both a decline in numbers of species recorded for order Rodentia (9 instead of 17 species) and slight increases in four orders Insectivora, Chiroptera, Primates and Pholidota (Table M2).

Table M2 Proportionate record of the known mammalian richness in Mabira FR in 2001 and 2006

Order	Known species number	% of known mammalian diversity recorded in 2001 study	% of known mammalian diversity recorded in 2006 study
Insectivora	6	33.3	50.0
Chiroptera	17	29.4	41.2
Primates	6	33.3	50.0
Carnivora	6	83.3	83.3
Artiodactyla	4	100.0	100.0
Pholidota	1	0.0	100.0
Hyracoidea	1	100.0	100.0
Rodentia	26	61.5	34.6
Macroscelidea	1	100.0	100.0

I have recorded just a little over 50% of the known mammal species for Mabira forest but slightly fewer species than were documented for the 2001 report.

Appendix M1 presents the species richness of mammals recorded in the different areas of the forest along the transmission line.

In total 35 species of mammals have been recorded for Mabira forest (Table M2) distributed in varying levels of species richness in the different transect locations. Although these results do not suggest that a total species inventory has been achieved, they however suggest some trends. The lowest species richness was recorded along transect 1 (Pylon 179) which, as observed from Section 2.121 was largely dominated by *Broussonetia papyrifera* and a very poor and sparse under storey. Transect 3 that was located in the forest along river Waliga returned the highest species richness of all sample locations in Mabira forest.




Figure M1. The cumulative number of species of small mammals (rodents & insectivores) recorded overall along the sample transects.

As is evident from Figure M1 a total species list was yet to be achieved since even after a cumulative sampling effort of 399 trap nights the graph has not reached an asymptote. The species recorded for these two taxa represent about 53% of those known to occur in Mabira forest implying that several more could have been recorded.

6.32 KIFU AND NAMYOYA FORESTS

Both these forests are very degraded, however for Kifu Forest Reserve the part to the south of the transmission line is in a secondary growth state with thick undergrowth among *Maesopsis eminii* (Section 2.122). The sections of the forest to the North of the transmission line are currently under plantation forests; the same is the case in Namyoya. The two are however under different management regimes with Kifu still under the direct control and management by the National Forestry Authority (NFA) while Namyoya was leased out to individual holders to grow trees. At the time we conducted the surveys for this report all the plots we visited in Namyoya had *Eucalyptus* growing or in some cases it had been harvested (Plates 12d, e & f).

The plantations are not very significant habitats for forest interior mammals because the complex undergrowth they depend on is lost in plantations (Plate 7). The undergrowth is important because it provides among other things: -

- (i) Cover for the mammals

- (ii) A source of food both for themselves and the other organisms they feed on.
- (iii) Maintains ambient environmental conditions for forest interior species

Table M3. Mammal species recorded from Kifu and Namyoya Central Forest Reserves

Mammal species	Kifu forest	Namyoya forest
<i>Crocidura olivieri</i> (Northern Giant Musk Shrew)	√	
<i>Crocidura turba</i> (Southern Woodland Musk Shrew)	√	
<i>Cercopithecus ascanius</i> (Red tailed Monkey) *	√	√
<i>Cercopithecus aethiops</i> (Vervet Monkeys)	√	√
<i>Cercoccebus albigena</i> (Grey Cheeked Mangabey) *	√	
<i>Potamochoerus porcus</i> (Bushpig)	√	
<i>Sylvicapra grimmia</i> (Common Duiker)	√	
<i>Canis adustus</i> (Side striped Jackal)	√	
<i>Civettictis citetta</i> (African Civet)	√	
<i>Panthera pardus</i> (Leopard)	√	
<i>Dendrohyrax arboreus</i> (Tree Hyrax) *	√	
<i>Thryonomys gregorianus</i> (Cane Rat)	√	
<i>Arvicanthis niloticus</i> (Nile rat)	√	√
<i>Dasymys incomtus</i> (Shaggy Marsh Rat)		√
<i>Grammomys dolichurus</i> (Common Thicket Rat)	√	
<i>Lemniscomys striatus</i> (Common striped Grass rat)	√	
<i>Lophuromys flavopunctatus</i> (Eastern Brush-furred Mouse)		√
<i>Lophuromys sikapusi</i> (Common Brush furred Mouse)	√	
<i>Mus minutoides</i> (Pygmy Mouse)		√
<i>Mus triton</i> (Grey-bellied Pygmy Mouse)	√	√
<i>Oenomys hypoxanthus</i> (Rusty nosed Rat)		√
<i>Xerus erythropus</i> (Stripped Ground Squirrel)		√
Totals	17	9

Altogether 22 species are presented in Table M4 for Kifu and Namyoya forests. Although Kifu CFR is badly degraded it still retains a fairly high species richness of mammals. However of these only three species (marked * Table M4), represent the forest interior species.

Namyoya forest on the other hand does not seem to retain much importance for forest interior mammals. The only such mammals that were recorded in this area are the Red tailed Monkeys. Given that the forest is now converted for growing *Eucalyptus* it is not likely that this species will survive in this area for very long. All other mammals recorded for this area are those of wide spread occurrence for which Namyoya is not a significant part of their range and/or for their conservation.

6.34 IMPORTANCE OF THE FOREST ALONG THE TRANSMISSION LINE FOR MAMMALS

The part of the Mabira forest that was surveyed for this report represents only a little over 1% of the total area of the Forest Reserve. The proportion of Mabira Forest's mammal species (50.7%) that have been documented in this small section of forest highlights its importance for mammal conservation within it.

Although some of the species recorded during these surveys (*A. paludinosus*, *M. longipes* and *D. ferrugineus*) appear in the IUCN records of assessed mammals (IUCN 2006), none of them is currently threatened. The majority of the species recorded are fairly common and widespread mammals except for the following, which are important regionally:

- 1 *Malacomys longipes* is a forest interior rodent with a niche strongly tied to riverine or other water-logged situations.
- 2 *Deomys ferrugineus* a strict forest interior species that never occurs in large densities
- 3 *Scutisorex somereni* a strict forest interior insectivore
- 4 *Rhynchocyon cirnei* a forest interior elephant shrew
- 5 *Rhinolophus alcyone* a forest interior microchiropteran
- 6 *Megaloglossus woermanni* a largely forest interior bat although it does extend its ranges into agro ecosystems typical of the Coffee/Banana systems.
- 7 *Crocidura selina* a Ugandan endemic shrew previously only known from Mabira Forest although it was subsequently recorded in other forests of Uganda

6.4 LUBIGI WETLANDS

The methods used here followed closely the approaches used in AESNP (2001) to enable acquisition of a primary data set that could be used in the future to assess potential impacts on the wetland ecosystem. Surveys in the wetlands were done for both any large and small mammals still present in the system.

The presence of larger mammals (Primates, Carnivores and Ungulates) was recorded through:

- (i) Direct observation for the diurnal species
- (ii) Indirect cues for their presence (such as spoors/paw/foot prints) and fecal pellets
- (iii) Recovery of skeletal material of species, or
- (iv) Interviews with local people.

The smaller mammals (rodents, insectivores and bats) were surveyed using traps.

Table M4 presents the record of mammal species recorded in the various sample sites of Lubigi wetlands.

A total of 16 species were recorded for the swamp, with 8 of these belonging to the single order Rodentia. Wetlands are usually not particularly rich in terrestrial biodiversity, but there is no reason to suppose that we recorded all possible mammals species in the swamp. An extended survey could record several other species. The seasonally flooding parts of the swamp could present suitable foraging and ranging areas for a variety of mammals.

In the swamp, primates are represented by the hardy Vervet monkey (*Cercopithecus aethiops*), which is only one of the very few primates that can still be found in human modified environments.

Owing to the location of Lubigi wetlands in a peri-urban setting and given the dense human population in areas surrounding the wetlands it is unlikely that Lubigi will be a significant ecosystem for much longer for larger mammals. From interviews with local people in Kazinga and Nganda Villages, Sitatunga and Bushbuck continue to be hunted for meat in the Lubigi wetlands. The wetlands are not receiving any active conservation, implying therefore that the surrounding local communities easily access and use resources in them.

Table M4. Mammal records for Lubigi Wetlands

	Sampling Sites along Lubigi Swamp							
	1	2	3	4	5	6	General Swamp records	
Insectivora								
Northern Giant Musk Shrew (<i>Crocidura olivieri</i>)	√							
Shrew (<i>Crocidura</i>)	√							
Primates								
Vervet Monkeys (<i>Cercopithecus eathiops</i>)	√			√	√		√	
Artiodactyla								
Bush Buck (<i>Tragelaphus scriptus</i>)		√						
Sitatunga (<i>Tragelaphus spekii</i>)	√	√	√	√	√	√	√	
Common Duiker (<i>Sylvicapra grimmia</i>)		√						
Carnivora								
Marsh Mongoose (<i>Atilax palludinosus</i>)	√		√	√	√		√	
Slender Mongoose (<i>Herpestes sanguineus</i>)		√		√	√	√	√	
Rodentia								
(<i>Aethomys kaiseri</i>)	√		√	√	√			
Nile Grass rat (<i>Arvicanthis niloticus</i>)	√		√	√				
Shaggy Swamp rat (<i>Dasymys incomtus</i>)	√	√	√	√				
Stripped Grass Mouse (<i>Lemniscomys striatus</i>)		√	√					
Brush-furred Mouse (<i>Lophuromys flavopunctatus</i>)	√	√	√	√	√	√		
Brush furred Mouse (<i>Lophuromys sikapusi</i>)	√	√	√	√	√	√		
Lesser cane Rat (<i>Thryonomys gregorianus</i>)	√		√		√			
Striped Ground Squirrel (<i>Xerus erythropus</i>)	√	√					√	

7.0 SUMMARY OF IMPACTS AND PROPOSED MITIGATION MEASURES

7.1 LIKELY IMPACTS

Table 7.1. Potential impacts in the forested areas

Possible effect / impact if not mitigated	Mitigation Options	Residual Impacts
1a. Forest cover (72 hectares) lost 1b. Habitat for under storey/forest interior species lost 1c. Forest edge to be extended deeper into forest 1d. Relatively intact forest to be lost between Pylons 148-170 1e. Forest fragmentation especially in Mabira CFR 2. Expansion of extent of the invasive <i>Broussonetia papyrifera</i> 3a. Rare species in direct impact zone lost 3b. Species richness and diversity lowered 4. Increased access into the forest possibly for illicit resource harvesting 5a. Habitat lost or degraded for riverine forest species 5b. Population fragmentation of forest interior understorey birds 5c. Behavioral disruption for mammals due to extended human presence 5d. Loss of breeding sites 5e. Loss of connectivity for understorey birds unwilling to cross the wayleave 6. Risk of collision by flying birds with conductors	1a. Provide compensation planted area of at least equivalent size to that to be cleared. 1b. Conduct enrichment planting with native plant species in Mabira and Kifu CFRs 1c. Allow for corridors between forest blocks north and south of transmission line 1d. Invest forest restoration in Kifu forest reserve 2. Conduct regular thinning out <i>B. papyrifera</i> or have it harvested by NFA and sold for fuel wood 3a. For plants carry out enrichment planting with in adjacent forest and the Kalagala offset. 3b. For animals, allow corridors of low vegetation to facilitate interconnectivity between forest sections. 4a. Strengthen forest Ranger outposts in areas close to the area of transmission line to provide enhanced policing of activities in the forest 4b. Increase community participation in managing Mabira and provide alternatives to reduce pressure on forest 5a. Preserve corridors of low vegetation along water courses crossing the wayleave 6. Attach reflectors to the conductors when finally installed	1a. Compensation area will not mirror exact forest state and biodiversity level to that lost 1b. Home ranges of forest interior/under storey species will be shrunk 1c. No overall loss of habitat if habitat creation is done early. 1d. Forest along the wayleave to have a richer influx of non forest species of for example butterflies 1e. Larval food plants and nectar plants are frequently found in large concentrations along roadways that similarly contain aggregations of pre- and -post diapause larvae of butterflies. 2. Invasive <i>B. papyrifera</i> could expand deeper into the natural forest. 3. No overall loss of biodiversity in Mabira forest if enrichment planting and restoration are done early 5a. No severe loss of habitat and biodiversity if corridors established early 5b. Reduced gene flow & consequently long term population viability 5c. Understorey takes decades to develop needing closure of canopy

Tables 7.1 and 7.2 summarise the main impacts, and the measures which might mitigate them to some extent. They are described in more detail in Section 7.3.

Plates 8 a – h and Plate 10 show a variety of impacts already happening on the swamp from actions of the neighbouring communities. Some of these actions such as harvesting papyrus and other grasses (Plate 8h), may be sustainable while mining for sand and clay for block making (Plates 8b and 10) alter the wetland considerably.

Table 7.2. Potential impacts in the wetland sites

Possible effect / impact if not mitigated	Mitigation Options	Residual Impacts
1a. Swamp Vegetation (~2 hectares) lost 1b. Habitat for typical swamp species lost	1a. Construct pylon footings out of or at edge of the swamp as much as is possible 1b. Discuss with Wetland Inspection Division options for better protection of the remaining intact swamp 1c. Increase community awareness and sensitization of the importance of wetlands	1. Swamp lost to infilling
2. Increased prominence of species not characteristic of swamp vegetation around pylons	2. Restrict the in filling to absolute minimum required for pylon footing	2. Swamp lost to infilling but no major loss of plant biodiversity in Lubigi Swamp
3a Species richness and diversity to be reduced 3b. Loss of breeding sites for animals	3a. Limit infilling and access routes into swamp to absolute minimum necessary	
4. Risk of collision by flying birds with conductors	4. Attach reflectors to the conductors when finally installed	

7.2 OVERALL IMPACT ASSESSMENT

For the different taxa we have covered for this assessment, many are widely distributed in Uganda in the right habitats with only a few that we have classified as rare because they were infrequently encountered in the transects or because they are actually known to be rare. The few Red Data species, as listed by IUCN, are unlikely to be seriously affected by the Bujagali Interconnection Project, and no specific mitigation measures are proposed for them. None of the potentially affected area is considered to be Critical Habitat as defined in IFC Performance Standard 6.

Except for the loss of a significant amount of forest and its attendant biodiversity it does not seem that the expansion of the wayleave for construction of the transmission line will result in major negative impacts on the terrestrial ecology. However, unless some forest vegetation, especially understorey vegetation, connects the north with the south of the forest, the negative effects of forest fragmentation will be increased.

With about 25 pylons installed along the wetland, this will translate into 1 ha of swamp lost and so will be its biodiversity and services. This increases the need for the remaining swamp to be better protected.

7.3 MANAGEMENT OPTIONS AND MITIGATION MEASURES

On the basis of our observations and the biological environment we have described in sections 2 – 6 we suggest the following as options for minimizing and mitigating the losses in biodiversity during and after the construction of the new transmission line and during maintenance activities along the wayleave.

1. Rare species (see Appendix P1) need special attention since they are the main contributors to diversity and conservation strategies should be laid out to protect them.
2. Mabira Forest is maintained in a pre-climax state by the anthropogenic pressure such as extraction of fuelwood, poles and even timber which if increased may retrogress the succession into a degraded community and, if decreased, may substitute *Broussonetia papyrifera* for other species such as *Celtis* spp and *Teclea nobilis*.
3. The southern side of the present powerline is relatively more degraded and more vulnerable to abuse because of the villages nearby, particularly near sites 7 and 8. It would, therefore, be best if the proposed transmission line was located on the southern side of the existing line, if that is possible.
4. There are various ways in which the forest lost to the wayleave could be compensated by improving the quality of other parts of the forest. FOREAIM is a project funded by the EU which is involved in restoration of degraded landscapes through a broad multidisciplinary approach. Using this approach, FOREAIM produces knowledge, practical tools, models and management guidelines for restoration implementation. With the full involvement of all stakeholders, FOREAIM synthesises information on economic, societal, policy and marketing issues to enhance employment opportunities and incomes, thus improving livelihoods for all sectors of the community. In Uganda, it is concentrating its efforts in Mabira and the progress is satisfactory. Given a history of degradation of Mabira forest, FOREAIM objectives are aimed at ensuring the natural regeneration of the forest, by considering the needs and expectations of the communities around. Support for this program, or at least adoption of its principles, would contribute to natural forest regeneration.
5. MAFICO a local Non-Governmental Organisation (NGO), is based at Najjembe, Mabira Forest Reserve. MAFICO aims to sensitise and educate communities on issues relating to the sustainable use of forest resources, ecotourism, agriculture, and income generating activities that support conservation. These would be useful allies for managing pressures on the forest.
6. The forest would benefit from enrichment-planting with high value timber species e.g. *Khaya anthotheca*, *Entandrophragma angolense*
7. Increased policing will be needed in Mabira since the expanded wayleave might give increased access into the forest.
8. The investor should replant a degraded forest area of equivalent or larger size to that which will be lost from Mabira, with indigenous trees.
9. Corridors of low vegetation should be maintained at the low points (Plates 4a & b) we have identified in the forest and vegetation along these should be left intact whenever the wayleave is cleared. These would enable understorey forest birds and other animals to maintain their link between the northern and southern sections of the forest. For this to be effective an “environmental management program” should be put in place for the Uganda Electricity Transmission Company (UETCL) where the supervisors will be able leave these locations,

and supervise the wayleave maintenance team accordingly.

10. If pylons in the Lubigi wetlands are to be placed at the same inter-pylon distance (about 400m), as is the case in Mabira forest, we envisage a total of about 25 pylons to be installed in the distance of about 10 km along the swamp. Given a footing of 0.04 ha for each pylon this will result into 1 ha of land offtake or wetland filling. As much as possible the pylons should be placed further landward to minimize the amount of swamp to be filled in.
11. The Lubigi swamp should be better protected. We were pleasantly surprised by the richness of the swamp flora and fauna, and the comparatively intact nature of large parts of the swamp. To judge from the EIA (Ministry of Works, Housing and Communications, 2001), the northern bypass to Kampala is likely to do far more damage to the swamp than the footings for the pylons. We propose that our report be drawn to the attention of the Wetlands Inspectorate Division in due course, with the recommendation that they give serious consideration to declaring as large a part as possible as some form of nature reserve. As well as the benefits to biodiversity and the environment generally, there are considerable educational possibilities for a site so close to Kampala.
12. Bird numbers in the three forests showed changes between the two years, but these may have been due to the data having been collected at different seasons and by different people. However, it is notable that there was also a drop in the numbers of forest mammal species, due mainly to there being far fewer rodents. These, like the forest birds, mainly inhabit the undergrowth; however, we observed no major changes in the vegetation. The reasons for the changes must therefore remain unexplained especially as there were no major changes observed in the vegetation. Continued monitoring of forest biodiversity is strongly recommended.
13. Improved management of Mabira Forest would probably more than compensate for the 72 hectares to be lost from widening the way leave. Illegal activities, in particular tree felling and trapping of mammals, appear to be common. But, as Naidoo and Adamowicz (2006) have shown, the potential income from ecotourists in Mabira could probably be increased considerably, for example by the ten-fold increase in fees that, they found, visitors would willingly pay and which, if used to improve forest management, would greatly benefit the conservation of the forest.

REFERENCES

- Beadle, L.C. and Lind, E. M. 1960. Research on the swamps of Uganda. *Uganda Journal* 24: 84 - 87.
- Behangana M. 1995: The Status of Amphibians in Rwenzori Mountains. MSc. Zoology Thesis, Makerere University, Kampala, Uganda.
- Bennun, L.A. and Njoroge, P. eds. 1996. Birds to watch in East Africa: a preliminary Red Data list. *Research Reports of the Centre for Biodiversity, National Museums of Kenya: Ornithology* 23: 1-16, Nairobi, Kenya.
- Bhatt, D. and Sanjit, L. 2005. How relevant are the concepts of species diversity and species richness? *Indian Academy of Sciences. Commentary; Journal of Bioscience* 30(5):101 – 000.
- Bibby, C.J.; N.J. Collar; M.J. Heath; C.H. Imboden; T.H. Johnston; A.J. Long, A.J. Stattersfiel and S.J. Thrgood, 1992. Putting Biodiversity on the map. Girton, Cambridge, UK.
- Blankenship, R.E. 2002. Molecular mechanisms of Photosynthesis. Blackwell Science, Oxford.
- Blauausten A.R. 1994: “Chicken or Nero’s Fiddle? A perspective on Declining Amphibian Populations”. *Herpetologica* 50: 85-57.
- Britton, P.L. 1978. Seasonality, density and diversity of birds of a papyrus swamp in western Kenya. *Ibis*, 120, 450-466.
- Brown, S. and Lugo, A.E. 1990. Tropical secondary forests. *Journal of Tropical Ecology* 6: 1–32.
- Byaruhanga, A., Kasoma, P. and Pomeroy, D. 2001. Important Bird Areas in Uganda. NatureUganda, Kampala, Uganda.
- Carswell, M. 1986. Birds of the Kampala area. *Scopus* special supplement Number 2 of the Ornithological Sub-committee of the East Africa Natural History Society, Nairobi, Kenya.
- Carswell, M., Pomeroy, D.E., Reynolds, J. and Tushabe, H. 2005. *Bird atlas of Uganda*. British Ornithologists’ Union, London.
- Carter, G. S. 1956. The papyrus swamps of Uganda. Heffer, Cambridge. England.
- Chazdon, R.L. 1998. Tropical forests – Log “em or leave “em? *Science* 281: 1295 – 1296.
- Colwell, R.K. 2001. Statistical estimation of species richness and shared species from samples, Version 6.0b1a. University of Connecticut. <http://viceroy.eeb.uconn.edu/estimate>.
- Colwell, R. K. and Coddington, J. A. 1994. Estimating terrestrial biodiversity through extrapolation. *Philosophical Transactions of the Royal Society (Series B)* 345: 101 – 118.
- Davenport T., Howard P. and Baltzer M. 1996. Mabira Forest Biodiversity Report. Report No. 13. Forest Department, Kampala, Uganda.
- De Witte, G.F. 1937. “Batraciens et Reptiles. Exploration du Parc Albert”. Mission, G..F. de Witte (1933-1935) *Inst. Parcs Nat. Congo Belge*, 33:xvii.
- Drewes, R.C. 1984. “A phylogenetic analysis of the Hyperoliidae (Anura): Tree frogs of Africa, Madagascar and the Seychelles. *Occasional Papers of the California Academic of Sciences*. 139, 1-70.
- Dranzoa, C. 1990. Survival of forest birds in formerly forested areas around Kampala. MSc thesis, Makerere University, Uganda.
- Drewes, R. C. & J. V. Vindum, 1994. “Amphibians of the Impenetrable Forest, South-Western Uganda”. *Journal of African Zoology* 108(1), 55-70.
- Dudley, C.O. 1978. The herpetofauna of the Lake Chilwa basin. *Nyala* 4: 87-99.
- Forest Department. 2002. Uganda Forestry Nature Conservation Master Plan. Ministry of Lands, Water and Environment, Kampala, Uganda.
- Freeman, N.S., Pomeroy, D.E. and Tushabe, H. 2003. On the use of Timed Species Counts to estimate abundance in species-rich communities. *African Journal of Ecology*, 41, 337-348.
- Gibbons, W.J. & D. R. Semlitsch, 1982: “Terrestrial Drift fences with Pitfalls for quantitative sampling of animal populations”. *Brimleyana*: No.7. 1-16.
- Guralnick, R. and Van Cleve, J. 2005. Strengths and weaknesses of museum and national survey data sets for predicting regional species richness: comparative and combined approaches. *Diversity and Distributions* 11: 349 – 359.
- Hamilton, A.C. 1982. Environmental history of East Africa. Academic Press, London, UK.
- Heltsche, G. F. and Forrester, J. H. 1983. Prediction of Neo-tropical tree and liana species richness from soil and climatic data. *Biod. Cons.* 4: 50 – 68.
- Heyer, W.R; Donnely, M.A.; McDiarmid, R.W.; Hayek Lee-Ann, C. AND Foster M.S. (eds) 1994: Measuring and monitoring Biological Diversity: Standard methods for Reptiles and Amphibians.

- Smithsonian Institution Press, PA, USA.
- Hobbs, R.J. 2000. Land use changes and invasions. In: Mooney, H.A. and Hobbs, R.J. (Eds.), *Invasive Species in a Changing World*. Island Press, Covelo, pp. 55 – 64.
- Howard, P.C. 1991. Conservation in Uganda's tropical Forest Reserves. IUCN, Gland, Switzerland and Cambridge, UK.
- International Union for Conservation of Nature (IUCN). 2000. Confirming the Global Extinction Crisis: A call for International Action as the most authoritative global assessment of species loss in released. <http://www.iucn.org/redlist/news.html>.
- IUCN 2006. 2006 IUCN Red List of Threatened Species. <www.iucnredlist.org>. Downloaded on 22 July 2006.
- Katende, A.B., Birnie, A and Tengnas, B. 1995. Useful trees and shrubs of Uganda. Regional Soil Conservation Unit/SIDA, *Technical Handbook Series* 10, Nairobi, Kenya.
- Katende, A.B., Birnie, A. and Tengnas, B. 2000. Useful trees and shrubs for Uganda. Identification, Propagation, and management for Agricultural and Pastoral communities. Regional Land Management Unit, SIDA, Nairobi.
- Khan, M.S. 1990. The imp-act of human activities on the status and distribution of amphibians in Pakistan. *Hamadryad*, 15: 21-24.
- Kigoolo, S. 1994: The impact of Wetland drainage and disturbance on the diversity of amphibians. M.Sc. Thesis, Zoology Department, Makerere University, Kampala, Uganda.
- Kovach, W. I. 1999. A multivariate statistical package for windows, version 3.1. Kovach computing services, Pentraeth, Wales.
- Kovach, W.I. 1999. A multivariate statistical package for windows, version 3.1. Kovach computing services, Pentraeth, Wales, UK.
- Langdale-Brown, I., Osmaston, H.A. & Wilson, G. 1964. The Vegetation of Uganda and its bearing on Land uses. Uganda Government Printer, Entebbe, Uganda.
- Loveridge, A. 1957: "Checklist of the reptiles and amphibians of East Africa". *Bulletin of the Museum of Comparative Zoology*, Harvard, Vol. 117(2) 153-362.
- Ludwig, J.A. and Reynolds, J.F. 1988. Statistical ecology: a primer on methods and computing. John Wiley and Sons, New York, USA.
- Mack, R.N., Simberloff, D., Lonsdale, W.M. 2000. Biotic invasions: causes, epidemiology, global consequences and control. *Ecological Applications* 10: 689-710.
- Maclean, M.D., Hassall, M., Boar, R.R. and Lake, I.R. 2006. Effects of disturbance and habitat loss on papyrus-dwelling passerines. *Biological Conservation* 131: 349-358.
- Mafabi, P. and Taylor, A.R.D. 1993. The National Wetlands Programme, Uganda In: Davis, T.J. 1993. Towards wise use of wetlands. Wise use project, Ramsar Convention Bureau, Gland, Switzerland.
- Maltby, E. 1991. Wetlands and their values. In: Finlayson, M. and Mosser, M 1991. Wetlands. International Waterfowl and Wetland Research Bureau (IWRB). Facts on File. Oxford, UK and NewYork, USA.
- McCollough et al, 1992: "Species Priorities and Methodological Constraints". Wildlife 2001. Populations. Elsevier Applied Science, London and New York, USA.
- Ministry of Natural Resources, 1992. Uganda Country Study on Costs, benefits and Unmet needs of Biological diversity conservation. Department of Environment Protection; National Biodiversity Unit, Kampala, Uganda.
- Ministry of Works, Housing and Communications, 2001. EIA – Kampala Northern Bypass. Entebbe, Uganda.
- Moss, B. 1980. Ecology of freshwaters: man and medium. Oxford Blackwell Scientific Publications, London, UK.
- Naidoo R. 2003. Economics, biodiversity conservation, and sustainable development. PhD thesis, Edmonton, University of Alberta, Canada.
- Naidoo, R. and Adamowicz, W.L., 2006. Economic benefits of biodiversity exceed costs of conservation at an African rainforest reserve. *Proceedings of the National Academy of Sciences* 102, 16,712-16,716.
- Namakambo N. 2000. District wetland inventory report Kampala. National wetlands programme WID, Kampala, Uganda
- National Environment Information Centre, 1992. Kampala District Profile. Ministry of Natural Resources, Kampala, Uganda.

- NBU, 1992. The Country's report of Country study on conservation of biodiversity in Uganda. Kampala, Uganda.
- Noble, G. K. 1954: The biology of the Amphibia. Dover Publications, New York, USA.
- Okland, R.H. 1990 Vegetation ecology: theory, methods and applications with reference to Fennoscandia. *Sommerfeltia supplement* 1: 4 – 233.
- Oliveira Filho, A.T., Mello, J.M., Scolforo, F.R.S. 1997. Effects of past disturbance and edges on tree community structure and dynamics within a fragment of tropical semideciduous forest in south-eastern Brazil over a five year period (1987 – 1992). *Plant Ecology* 131: 45 – 66.
- Phillips, K. 1990. Where have all the frogs and toads gone?. *BioScience* 40: 422-424.
- Polhill, R.M., Milne-Redhead, E., Turrill, W.B., and Hubbard, C.E. (eds) (from 1952 onwards). Flora of Tropical East Africa (in many parts). Crown Agents, London and Balkema, Rotterdam, Holland.
- Pomeroy D.E, Kityo R.M. & Ssegawa P. 2006 Environment Impact Assessment study for Bujagali Hydropower Proposed Dam site (Terrestrial Biodiversity component).
- Rossouw J & Sacchi M 1998. Where to watch birds in Uganda UTB, Kampala, Uganda.
- Saldarriaga, J.G. and Uhl, C. 1991. Recovery of forest vegetation following slash-and-burn agriculture in the Upper Rio Negro. In: Tropical Rainforests: regeneration and management (A. Gomez-Pompa, T.C. Whitmore, and M. Hadley eds.) Blackwell, New York, USA. p. 303 – 312.
- Schiotz, A. 1972(a): "The Superspecies *Hyperolius viridiflavus* (Anura)". *Vidensk. Meddr dansk naturh. Foren.* 134:21-76.
- Schiotz, A. 1972(b): "The Treefrogs of Eastern Africa". *Spolia zool. Mus. Haun.* 25: 1-346.
- Sharma, G.P. & Raghubanshi, A. S. 2005. Impact of *Lantana camara* L. invasion on herb layer diversity and soil properties in a dry deciduous forest. *Journal of Ecology and Application* 11(1): 27 – 39.
- Tilman, D. 1982. Resource competition and community structure. Princeton University Press.
- U.S. EPA. 2002. Methods for Evaluating Wetland Condition: Using Amphibians in Bioassessments of Wetlands. Office of Wetlands, U.S. Environmental Protection Agency, Washington, DC, USA. EPA-822-R-02-022.
- Visser, A. 1960. A study of the decomposition of papyrus in the swamps of Uganda. *PhD Thesis*. University of London, UK
- W.C.M.C. 1991: Animals of Uganda, Conservation status listings. Cambridge, UK.
- Welch K.R.C. 1982. Herpetology of Africa. A Checklist and bibliography of the orders amphibians, sauria and serpents.
- Wettstein, W and Schmid, B 1999. Conservation of Arthropod diversity in montane wetlands: effect of altitude, habitat quality and habitat fragmentation on butterflies and grasshoppers. *Journal of Applied Ecology* 1999, 36: 363 – 373
- Young, J. Z. 1981: The life of vertebrates. 3rd ed. Oxford University Press, London, UK.

Appendix F1. Figures (Maps)

- Figure 1. Mabira Location Plan***
Figure 2. Lubigi Location Plan
Figure 3. Lubigi Sampling Site Locations
Figure 4. Dominant Plant Formations (Lubigi)
Figure 5. Human Impacts (Lubigi)

Figure 1. The Mabira Forest Reserve Environs showing Sites where Detailed Ecological Surveys were Conducted




Figure 2. Lubigi Location Plan




Figure 3: Sampling Site Locations on 1:50,000 Topomap Base (Lands and Surveys Department, 1969)




Figure 4: Dominant Plant Formation(s)





Figure 5: Human Impacts



Appendix G1. GPS locations of landmarks that were noted in Mabira, Kifu and Namyoya Forests

Land Mark	Latitude	Longitude	Altitude (m)
Mabira Forest readings			
Site 1 – Pylon 179	N00°27.1202'	E033°06.1618'	1216.9
Site 2 – Pylon 174	N00°27.0178'	E033°05.3587'	1173.2
Site 3 – Pylon 170/169	N00°26.9230'	E033°04.5116'	1141.4
Site 4 – Pylon 164	N00°26.7980'	E033°03.4528'	1153.2
Site 5 – Pylon 158	N00°26.5470'	E033°01.7723'	1176.8
Site 6 – Pylon 154	N00°26.4410'	E033°00.8432'	1154.9
Site 7 – Pylon 149	N00°26.6517'	E033°02.4092'	1204.4
Site 8 – Pylon 144	N00°26.3183'	E032°59.9755'	1167.4
Cultivation along wayleave	N00°26.2285'	E032°59.3487'	1215.0
Stream with water	N00°26.5123'	E033°01.3877'	1159.5
Stream with water	N00°26.5012'	E033°01.2997'	1147.2
Stream with water	N00°26.4943'	E033°01.2399'	1154.9
Stream with water	N00°26.3682'	E033°00.3463'	1157.5
Stream with water	N00°26.3469'	E033°00.2221'	1158.0
Stream with water	N00°26.3424'	E033°00.1773'	1166.0
Valley point at a steep drop near Pylon 141	N00°26.2701'	E032°59.6397'	1256.6
Temporary Pond	N00°26.6133'	E033°02.0754'	1189.7
Kifu Forest readings			
Pylon 64	N00°23.928'	E032°45.254'	
<i>Auracaria</i> Plantation	N00°23.954'	E032°45.359'	
Namyoya Forest readings			
Pylon 51	N00°23.611'	E032°42.749'	
Harvested <i>Eucalyptus</i> plot	N00°23.634'	E032°42.894'	
Wetland	N00°23.637'	E032°42.989'	

Appendix G2. GPS locations of landmarks that were noted along the western border of Lubigi Swamp

Land Mark	Latitude	Longitude	Altitude (m)
	N00°19.2496'	E032°30.8257'	1159.0
	N00°19.2755'	E032°30.8276'	1159.2
	N00°19.2963'	E032°30.8475'	1159.0
	N00°19.3067'	E032°30.8233'	1160.2
	N00°19.3279'	E032°30.8156'	1160.2
	N00°19.3549'	E032°30.8954'	1155.9
	N00°19.3809'	E032°30.8646'	1162.1
	N00°19.4165'	E032°30.8654'	1160.2
	N00°19.4126'	E032°30.8908'	1159.7
	N00°20.1655'	E032°31.2236'	1163.1
	N00°18.1515'	E032°30.4714'	1162.6
	N00°18.3314'	E032°30.4273'	1161.6
	N00°18.3509'	E032°30.4222'	1163.1
	N00°18.0342'	E032°30.8065'	1162.6
	N00°18.0234'	E032°30.9017'	1155.6
Swamp Site 4 (Lubanyi 2)	N00°19.3867'	E032°30.8364'	1161.4
	N00°18.0544'	E032°30.6420'	1163.1
	N00°20.1431'	E032°31.2487'	1165.2
	N00°18.0439'	E032°30.7200'	1162.6
	N00°20.1307'	E032°31.2723'	1164.8
	N00°18.1879'	E032°30.4494'	1162.6
	N00°18.0420'	E032°30.7499'	1158.0
	N00°18.1223'	E032°30.4918'	1160.7
Swamp Site 6 (Ganda)	N00°20.7580'	E032°31.9657'	1170.8
Swamp Site 2 (Kazinga 2)	N00°18.1497'	E032°30.4729'	1162.4
	N00°18.2754'	E032°30.4233'	1163.8
	N00°18.0399'	E032°30.7679'	1163.3
	N00°18.0321'	E032°30.8443'	1159.9
	N00°18.9849'	E032°30.7235'	1160.4
	N00°20.0647'	E032°31.3053'	1168.6
	N00°20.9327'	E032°32.3873'	1168.8
Swamp Site 1 (Kazinga 1)	N00°18.0655'	E032°30.5876'	1160.4
	N00°18.0878'	E032°30.5172'	1161.9
Swamp Site 3 (Lubanyi 1)	N00°19.2612'	E032°30.7911'	1159.7
Swamp Site 5 (Nakawudde)	N00°20.2028'	E032°31.2075'	1165.2
Rubish dump	N00°18.075'	E032°30.545'	
Brick Making	N00°18.082'	E032°30.520'	
Sewage	N00°18.101'	E032°30.417'	
Grass harvesting	N00°18.101'	E032°30.417'	
Potato Garden	N00°18.108'	E032°30.388'	
Old tyres burnt	N00°18.122'	E032°30.348'	
Block making	N00°18.130'	E032°30.301'	

Appendix P 1: Plant species recorded from Mabira, Kifu and Namyoya forest reserves

1 means presence of species at site

H = Herb; C = Climber; T = Tree, S = Shrub

Abundance: D – Dominant, A –Abundant, F – Frequent, O – Occasional, R - Rare

Family	Species	Life form	Abundance	Mabira F.R. Sites								Kifu F.R.	Namyoya F.R.	Totals
				1	2	3	4	5	6	7	8	9	10	
Acanthaceae	<i>Acanthus pubescens</i>	S	O									1	1	2
Acanthaceae	<i>Asystasia gigantea</i>	H	O	1							1			2
Acanthaceae	<i>Dicliptera laxata</i>	H	R	1										1
Acanthaceae	<i>Dyschoriste radicans</i>	H	O		1	1								2
Acanthaceae	<i>Hypoestes sp.</i>	H	R									1		1
Acanthaceae	<i>Justicia flava</i>	H	O		1							1		2
Acanthaceae	<i>Justicia scandens</i>	H	O	1		1								2
Acanthaceae	<i>Justicia sp.</i>	H	R						1					1
Acanthaceae	<i>Justicia striata</i>	H	R							1				1
Acanthaceae	<i>Lankasteria elegans</i>	H	R		1									1
Acanthaceae	<i>Phaulopsis angolana</i>	H	R									1		1
Acanthaceae	<i>Pseuderanthemum ludovicianum</i>	H	F			1	1	1		1	1			5
Alangiaceae	<i>Alangium chinense</i>	T	O						1	1				2
Amaranthaceae	<i>Achyranthes aspera</i>	H	A		1	1		1	1	1	1	1		7
Amaranthaceae	<i>Aerva lanata</i>	H	R			1								1
Amaranthaceae	<i>Amaranthus spinosa</i>	H	O				1			1				2
Amaranthaceae	<i>Psilotricum scleranthum</i>	H	R				1							1
Anacardiaceae	<i>Pseudospondias microcarpa</i>	T	F				1		1	1	1	1		5
Annonaceae	<i>Monodora myrsitica</i>	T	R						1					1
Annonaceae	<i>Uvariopsis congensis</i>	C	R						1					1
Apocynaceae	<i>Alafia caudata</i>	C	R				1							1
Apocynaceae	<i>Alafia scandens</i>	C	O	1			1							2
Apocynaceae	<i>Alstonia boonei</i>	T	F			1	1	1		1	1			5
Apocynaceae	<i>Funtumia africana</i>	T	F					1	1		1			3
Apocynaceae	<i>Funtumia elastica</i>	T	A		1	1		1	1	1	1	1	1	8
Apocynaceae	<i>Motandra guineensis</i>	C	R			1								1
Apocynaceae	<i>Rauvolfia caffra</i>	T	R			1								1
Apocynaceae	<i>Tabernamontana holstii</i>	T	O					1				1		2
Araceae	<i>Culcasia fulcifolia</i>	H	O		1	1								2

Appendix P1 (Continued)

Family	Species	Life form	Abundance	Mabira F.R. Sites								Kifu F.R.	Namyoya F.R.	Totals
				1	2	3	4	5	6	7	8	9	10	
Araceae	<i>Culcasia scandens</i>	H	F					1	1	1	1	1		5
Aristolochiaceae	<i>Aristolochia elegans</i>	C	R					1						1
Aristolochiaceae	<i>Aristolochia triactina</i>	C	R	1										1
Asclepiadaceae	<i>Mondia whytei</i>	C	R		1									1
Asclepiadaceae	<i>Pentarrhinum abyssinicum</i>	C	R		1									1
Asclepiadaceae	<i>Secamone africana</i>	C	R									1		1
Asclepiadaceae	<i>Secamone sp.</i>	C	R		1									1
Aspleniaceae	<i>Asplenium sp.</i>	H	R			1								1
Asteraceae	<i>Acmella caulorrhiza</i>	H	R				1							1
Asteraceae	<i>Ageratum conyzoides</i>	H	A			1	1			1	1	1	1	6
Asteraceae	<i>Aspilia africana</i>	H	O	1			1							2
Asteraceae	<i>Bidens pilosa</i>	H	F		1	1						1	1	4
Asteraceae	<i>Conyza floribunda</i>	H	A	1	1				1	1	1		1	6
Asteraceae	<i>Crassocephalum montuosum</i>	H	R										1	1
Asteraceae	<i>Melanthera scandens</i>	H	O				1					1		2
Asteraceae	<i>Senecio syringifolius</i>	C	R								1			1
Asteraceae	<i>Synedrella nodiflora</i>	H	A		1	1	1			1	1	1	1	7
Asteraceae	<i>Tithonia diversifolia</i>	S	O							1	1			2
Asteraceae	<i>Vernonia amygdalina</i>	S	O									1	1	2
Araucariaceae	<i>Araucaria cunninghamii</i>	T	R									1		1
Balanitaceae	<i>Balanites wilsoniana</i>	T	O			1	1							2
Bignoniaceae	<i>Kigelia africana</i>	T	R		1									1
Bignoniaceae	<i>Markhamia lutea</i>	T	A	1		1	1	1	1	1	1	1		8
Bignoniaceae	<i>Spathodea campanulata</i>	T	R				1							1
Burseraceae	<i>Canarium schweinfurthii</i>	T	O				1		1					2
Capparidaceae	<i>Capparis erythrocarpos</i>	C	R		1									1
Capparidaceae	<i>Capparis tomentosa</i>	C	R	1										1
Capparidaceae	<i>Maerua duchesnei</i>	S	O				1				1			2
Capparidaceae	<i>Ritchiea afzeli</i>	T	O				1		1					2
Cecropiaceae	<i>Myrianthus arboreus</i>	T	R								1			1
Celastraceae	<i>Pristimera plumbea</i>	C	R									1		1
Celastraceae	<i>Salacia leptoclada</i>	C	O				1				1			2
Combretaceae	<i>Combretum paniculatum</i>	C	R						1					1

Appendix P1 (Continued)

Family	Species	Life form	Abundance	Mabira F.R. Sites								Kifu F.R.	Namyoya F.R.	Totals
				1	2	3	4	5	6	7	8	9	10	
Commelinaceae	<i>Commelina africana</i>	H	O		1	1								2
Commelinaceae	<i>Commelina benghalensis</i>	H	R									1		1
Commelinaceae	<i>Commelina latifolia</i>	H	F					1	1	1	1			4
Commelinaceae	<i>Palisota manii</i>	H	O			1						1		2
Commelinaceae	<i>Pollia condensata</i>	H	A		1	1	1		1	1	1			6
Connaraceae	<i>Agelaea pentagyna</i>	C	R			1								1
Connaraceae	<i>Agelaea ugandensis</i>	C	F			1			1		1			3
Connaraceae	<i>Cnestis ugandensis</i>	S	R							1				1
Connaraceae	<i>Connaras longistipitatus</i>	C	R								1			1
Convolvulaceae	<i>Dichondra repens</i>	H	R							1				1
Convolvulaceae	<i>Evolvulus nummularius</i>	H	F			1		1	1	1	1			5
Convolvulaceae	<i>Hewittia sublobata</i>	C	F	1								1	1	3
Convolvulaceae	<i>Ipomoea acuminata</i>	C	R			1								1
Convolvulaceae	<i>Ipomoea batatus</i>	C	R										1	1
Convolvulaceae	<i>Ipomoea grantii</i>	C	R									1		1
Convolvulaceae	<i>Lepistemon owariense</i>	C	R									1		1
Convolvulaceae	<i>Stictocardia beraviensis</i>	C	R									1		1
Cucurbitaceae	<i>Cucumis figarei</i>	C	R			1								1
Cucurbitaceae	<i>Momordica foetida</i>	C	F		1	1			1		1	1		5
Cucurbitaceae	<i>Mukia maderaspatana</i>	C	R					1						1
Cyperaceae	<i>Cyperus cyperoides</i>	H	R										1	1
Cyperaceae	<i>Cyperus distans</i>	H	O									1	1	2
Cyperaceae	<i>Fimbristylis dichotoma</i>	H	R				1							1
Cyperaceae	<i>Kyllinga aurata</i>	H	O			1						1		2
Davalliaceae	<i>Arthropteris orientalis</i>	H	O					1			1			2
Davalliaceae	<i>Arthropteris palisoti</i>	H	R			1								1
Dichapetalaceae	<i>Tapura fischeri</i>	S	F		1	1				1				3
Dilleniaceae	<i>Tetracera potatoria</i>	C	F		1	1		1	1					4
Dioscoreaceae	<i>Dioscorea abyssinica</i>	C	R			1								1
Dracaenaceae	<i>Dracaena fragrans</i>	S	A	1	1	1	1	1	1	1	1			8
Dracaenaceae	<i>Dracaena laxissima</i>	C	R				1							1
Ebenaceae	<i>Diospyros abyssinica</i>	T	F	1				1	1					3
Euphorbiaceae	<i>Acalypha acrogyna</i>	S	R	1										1

Appendix P1 (Continued)

Family	Species	Life form	Abundance	Mabira F.R. Sites								Kifu F.R.	Namyoya F.R.	Totals
				1	2	3	4	5	6	7	8	9	10	
Euphorbiaceae	<i>Acalypha bipartita</i>	S	A	1	1	1		1	1	1		1		7
Euphorbiaceae	<i>Acalypha neptunica</i>	S	D	1	1	1	1	1	1	1	1	1		9
Euphorbiaceae	<i>Acalypha ornata</i>	S	F			1				1		1	1	4
Euphorbiaceae	<i>Alchornea cordifolia</i>	T	O									1	1	2
Euphorbiaceae	<i>Argomuellera macrophylla</i>	S	A	1	1	1	1	1	1	1	1			8
Euphorbiaceae	<i>Bridelia micrantha</i>	T	F	1						1		1		3
Euphorbiaceae	<i>Croton macrostachyus</i>	T	A			1	1	1	1	1		1		6
Euphorbiaceae	<i>Croton megalocarpus</i>	T	R					1						1
Euphorbiaceae	<i>Erythrococca sp.</i>	S	R									1		1
Euphorbiaceae	<i>Erythrococca stolziana</i>	S	F	1	1	1	1							4
Euphorbiaceae	<i>Euphorbia hirta</i>	H	R			1								1
Euphorbiaceae	<i>Mallotus oppositifolius</i>	S	R			1								1
Euphorbiaceae	<i>Manhot esculentus</i>	S	R										1	1
Euphorbiaceae	<i>Margaritaria discoides</i>	T	F			1	1	1		1	1			5
Euphorbiaceae	<i>Neobotonia melleri</i>	T	R							1				1
Euphorbiaceae	<i>Phyllanthus amarus</i>	H	F	1	1								1	3
Euphorbiaceae	<i>Phyllanthus ovalifolius</i>	S	R									1		1
Euphorbiaceae	<i>Ricinus communis</i>	S	O	1	1									2
Euphorbiaceae	<i>Sapium ellipticum</i>	T	A	1			1	1	1			1	1	6
Euphorbiaceae	<i>Spondianthus preusii</i>	T	R									1		1
Euphorbiaceae	<i>Tragia brevipes</i>	C	O			1							1	2
Fabaceae	<i>Acacia pentagona</i>	C	A			1		1	1	1	1	1		6
Fabaceae	<i>Albizia coriaria</i>	T	O							1	1			2
Fabaceae	<i>Albizia glaberrima</i>	T	A	1		1	1	1	1	1	1			7
Fabaceae	<i>Albizia grandibracteata</i>	T	A	1			1	1	1			1	1	6
Fabaceae	<i>Albizia gummifera</i>	T	A	1		1	1	1	1	1	1			7
Fabaceae	<i>Albizia zygia</i>	T	F							1	1	1		3
Fabaceae	<i>Baikiaea insignis</i>	T	F			1	1		1					3
Fabaceae	<i>Baphiopsis parviflora</i>	T	R						1					1
Fabaceae	<i>Cassia spectabilis</i>	T	R										1	1
Fabaceae	<i>Centrosema pubescens</i>	C	O			1							1	2
Fabaceae	<i>Crotalaria sp.</i>	S	O					1	1					2
Fabaceae	<i>Dalbergia lactea</i>	C	O							1		1		2

Appendix P1 (Continued)

Family	Species	Life form	Abundance	Mabira F.R. Sites								Kifu F.R.	Namyoya F.R.	Totals
				1	2	3	4	5	6	7	8	9	10	
Fabaceae	<i>Desmodium adscendens</i>	H	F	1			1					1	1	4
Fabaceae	<i>Desmodium repandum</i>	H	A	1	1		1	1	1	1	1	1		8
Fabaceae	<i>Desmodium salicifolium</i>	H	R			1								1
Fabaceae	<i>Desmodium triflorum</i>	H	O				1	1						2
Fabaceae	<i>Desmodium velutinum</i>	S	R	1										1
Fabaceae	<i>Erythrina abyssinica</i>	T	R			1								1
Fabaceae	<i>Glycine wightii</i>	H	A	1	1		1	1	1	1		1		7
Fabaceae	<i>Indigofera spicata</i>	H	A	1	1	1	1	1	1					6
Fabaceae	<i>Mimosa pudica</i>	H	A			1	1	1	1	1	1			6
Fabaceae	<i>Parkia filicoidea</i>	T	R									1		1
Fabaceae	<i>Peptadeniastrum africanum</i>	T	F	1	1			1						3
Fabaceae	<i>Rhynchosia sublobata</i>	C	R									1		1
Fabaceae	<i>Senna hirsuta</i>	S	D	1	1		1	1	1	1	1	1	1	9
Fabaceae	<i>Teramnus labialis</i>	C	A	1		1	1	1	1	1				6
Fabaceae	<i>Vigna unguiculata</i>	H	R		1									1
Flacourtiaceae	<i>Dovyalis macrocalyx</i>	S	R				1							1
Flacourtiaceae	<i>Lindackeria lanceolata</i>	S	R		1									1
Guttiferae	<i>Harungana madagascariensis</i>	T	O									1	1	2
Hernandiaceae	<i>Illigera pentaphylla</i>	C	O		1						1			2
Labiatae	<i>Leonotis nepetifolia</i>	S	R										1	1
Labiatae	<i>Leucas martinicensis</i>	H	R			1								1
Labiatae	<i>Ocimum gratissimum</i>	S	A	1	1	1	1		1	1	1			7
Labiatae	<i>Stachys argillicola</i>	H	F				1	1	1	1	1			5
Leeaceae	<i>Leea guineensis</i>	S	O			1						1		2
Malpighiaceae	<i>Flabellaria paniculata</i>	C	R				1							1
Malvaceae	<i>Hibiscus calyphyllus</i>	H	F	1								1	1	3
Malvaceae	<i>Hibiscus surrantensis</i>	H	R									1		1
Malvaceae	<i>Pavonia urens</i>	S	R	1										1
Malvaceae	<i>Sida acuta</i>	H	A		1		1	1	1	1	1	1	1	8
Malvaceae	<i>Sida rhombifolia</i>	H	A		1	1	1	1	1	1				6
Malvaceae	<i>Urena lobata</i>	H	A	1	1	1	1					1	1	6
Marantaceae	<i>Marantochloa leucantha</i>	H	F			1			1	1		1		4
Melastomataceae	<i>Memecylon jasminoides</i>	S	R		1									1

Appendix P1 (Continued)

Family	Species	Life form	Abundance	Mabira F.R. Sites								Kifu F.R.	Namyoya F.R.	Totals
				1	2	3	4	5	6	7	8	9	10	
Meliaceae	<i>Entandrophragma angolense</i>	T	O						1		1			2
Meliaceae	<i>Entandrophragma cylindricum</i>	T	R						1					1
Meliaceae	<i>Khaya anthotheca</i>	T	O	1						1				2
Meliaceae	<i>Trichilia dregeana</i>	T	F					1	1	1				3
Meliaceae	<i>Trichilia fischeri</i>	T	F				1	1	1					3
Meliaceae	<i>Trichilia preuriana</i>	T	A			1	1	1	1	1	1			6
Meliaceae	<i>Trichilia rubescens</i>	T	R						1					1
Meliaceae	<i>Turraea vogellioides</i>	T	R				1							1
Menispermaceae	<i>Cissampelos mucronata</i>	C	F					1	1				1	3
Menispermaceae	<i>Tinospora caffra</i>	C	R					1						1
Moraceae	<i>Antiaris toxicaria</i>	T	A		1	1		1	1	1	1	1	1	8
Moraceae	<i>Artocarpus heterophyllus</i>	T	F	1	1								1	3
Moraceae	<i>Broussonetia papyrifera</i>	T	A	1	1	1	1	1	1	1	1			8
Moraceae	<i>Ficus asperifolia</i>	T	F	1				1				1		3
Moraceae	<i>Ficus brachylepis</i>	T	R				1							1
Moraceae	<i>Ficus exasperata</i>	T	A	1	1		1	1	1	1	1		1	8
Moraceae	<i>Ficus lingua</i>	T	R							1				1
Moraceae	<i>Ficus mucoso</i>	T	F				1	1		1				3
Moraceae	<i>Ficus ovata</i>	T	R			1								1
Moraceae	<i>Ficus polita</i>	T	F			1	1		1		1			4
Moraceae	<i>Ficus sur</i>	T	F			1	1				1	1	1	5
Moraceae	<i>Ficus vallis-chaude</i>	T	R							1				1
Moraceae	<i>Mesozygia lactea</i>	T	R			1								1
Moraceae	<i>Morus mesozygia</i>	T	R						1					1
Moraceae	<i>Trilepisium madagascariensis</i>	T	A	1		1	1	1	1	1	1	1		8
Musaceae	<i>Musa sapientum</i>	H	R						1					1
Myrsinaceae	<i>Ardisia staudtii</i>	S	R							1				1
Myrsinaceae	<i>Maesa lanceolata</i>	T	R									1		1
Myrsinaceae	<i>Maesa welwitschii</i>	C	R		1									1
Myrtaceae	<i>Eucalyptus grandis</i>	T	R										1	1
Myrtaceae	<i>Eugenia emens</i>	S	O					1	1					2
Myrtaceae	<i>Psidium guajava</i>	T	O	1									1	2
Myrtaceae	<i>Syzygium guineense</i>	T	R								1			1

Appendix P1 (Continued)

Family	Species	Life form	Abundance	Mabira F.R. Sites								Kifu F.R.	Namyoya F.R.	Totals
				1	2	3	4	5	6	7	8	9	10	
Nyctaginaceae	<i>Pisonia aculeata</i>	C	R							1				1
Ochanaceae	<i>Ouratea densiflora</i>	T	R					1						1
Oleaceae	<i>Jasminum eminii</i>	C	O	1						1				2
Oleandraceae	<i>Schrebera arborea</i>	T	R								1			1
Oxalidaceae	<i>Oxalis corniculata</i>	H	R							1				1
Passifloraceae	<i>Adenia aculeata</i>	C	R					1						1
Passifloraceae	<i>Adenia cissampeloides</i>	C	O					1		1				2
Passifloraceae	<i>Passiflora edulis</i>	C	R		1									1
Phytolaccaceae	<i>Phytolacca dodecandra</i>	C	O		1					1				2
Piperaceae	<i>Piper umbellatum</i>	C	F	1		1					1	1		4
Plumbaginaceae	<i>Plumbago zeylanica</i>	H	R			1								1
Poaceae	<i>Acroceras zizanioides</i>	Gr	R									1		1
Poaceae	<i>Brachiaria decumbens</i>	Gr	R										1	1
Poaceae	<i>Chloris pycnothrix</i>	Gr	O	1	1									2
Poaceae	<i>Cynodon aethiopicus</i>	Gr	O		1	1								2
Poaceae	<i>Cynodon dactylon</i>	Gr	F	1			1	1	1	1				5
Poaceae	<i>Digitaria abyssinica</i>	Gr	R										1	1
Poaceae	<i>Eleusine indica</i>	Gr	O		1		1							2
Poaceae	<i>Eragrostis tunuifolia</i>	Gr	R		1									1
Poaceae	<i>Isachne buettneri</i>	Gr	R									1		1
Poaceae	<i>Leptaspis cochleata</i>	Gr	F			1		1	1					3
Poaceae	<i>Olyra latifolia</i>	Gr	R							1				1
Poaceae	<i>Oplismenus hirtellus</i>	Gr	F	1	1							1		3
Poaceae	<i>Panicum maximum</i>	Gr	F	1	1	1							1	4
Poaceae	<i>Panicum trichocladum</i>	Gr	F	1								1	1	3
Poaceae	<i>Panicum vaginatum</i>	Gr	R									1		1
Poaceae	<i>Paspalum conjugatum</i>	Gr	A	1	1	1	1	1	1	1		1		8
Poaceae	<i>Paspalum scrobiculatum</i>	Gr	O		1							1		2
Poaceae	<i>Pennisetum polystachion</i>	Gr	R		1									1
Poaceae	<i>Pennisetum purpureum</i>	Gr	R		1									1
Poaceae	<i>Pseudobromus silvaticus</i>	Gr	R								1			1
Poaceae	<i>Rottboelia conchinchinensis</i>	Gr	R										1	1
Poaceae	<i>Setaria megaphylla</i>	Gr	O	1								1		2

Appendix P1 (Continued)

Family	Species	Life form	Abundance	Mabira F.R. Sites								Kifu F.R.	Namyoya F.R.	Totals
				1	2	3	4	5	6	7	8	9	10	
Poaceae	<i>Sporobolus africanus</i>	Gr	R	1										1
Poaceae	<i>Sporobolus pyramidalis</i>	Gr	R		1									1
Pteridaceae	<i>Pteris dentata</i>	H	O							1		1		2
Rhamnaceae	<i>Gouania longispicata</i>	T	R									1		1
Rhamnaceae	<i>Maesopsis eminii</i>	T	F	1		1	1					1		4
Rhamnaceae	<i>Scutia myrtina</i>	S	F			1	1	1				1		4
Rhamnaceae	<i>Ventilago diffusa</i>	C	R			1								1
Rosaceae	<i>Rubus apetalus</i>	S	O	1								1		2
Rubiaceae	<i>Canthium lactescens</i>	T	O		1			1						2
Rubiaceae	<i>Coffea canephora</i>	S	O						1		1			2
Rubiaceae	<i>Dictyandra arborescens</i>	S	R						1					1
Rubiaceae	<i>Geophila repens</i>	H	O		1						1			2
Rubiaceae	<i>Hymenocoleus hirsuta</i>	H	R				1							1
Rubiaceae	<i>Oxyanthus subpunctatus</i>	S	R		1									1
Rubiaceae	<i>Psychotria sp.</i>	S	R									1		1
Rubiaceae	<i>Spermacoce princeae</i>	H	R									1		1
Rubiaceae	<i>Uncaria africana</i>	C	R				1							1
Rutaceae	<i>Chaetachme aristata</i>	S	F			1			1			1		3
Rutaceae	<i>Citropsis articulata</i>	S	F	1		1		1		1	1			5
Rutaceae	<i>Clausena anisata</i>	S	O				1					1		2
Rutaceae	<i>Fagaropsis angolensis</i>	T	O					1	1					2
Rutaceae	<i>Rothmannia urcelliformis</i>	T	R			1								1
Rutaceae	<i>Rutidea orientalis</i>	C	R				1							1
Rutaceae	<i>Teclea nobilis</i>	T	A	1		1	1	1	1	1		1		7
Rutaceae	<i>Toddalia asiatica</i>	S	F				1		1			1		3
Rutaceae	<i>Zanthoxylum gillettii</i>	T	F				1		1		1			3
Sapindaceae	<i>Allophylus africana</i>	S	O	1								1		2
Sapindaceae	<i>Allophylus macrobotrys</i>	S	R				1							1
Sapindaceae	<i>Aphania senegalensis</i>	T	F			1	1	1	1	1				5
Sapindaceae	<i>Blighia unijugata</i>	T	F			1	1	1	1		1			5
Sapindaceae	<i>Cardiospermum grandiflorum</i>	C	O			1						1		2
Sapindaceae	<i>Cardiospermum halicacabum</i>	C	F	1		1	1					1		4
Sapindaceae	<i>Deinbollia kilimandscharica</i>	T	R			1								1

Appendix P1 (Continued)

Family	Species	Life form	Abundance	Mabira F.R. Sites								Kifu F.R.	Namyoya F.R.	Totals
				1	2	3	4	5	6	7	8	9	10	
Sapindaceae	<i>Glenniea africana</i>	T	F			1		1	1					3
Sapindaceae	<i>Lasciodiscus mildbraedii</i>	T	A		1	1	1	1	1	1	1			7
Sapindaceae	<i>Lychnodiscus cerospermus</i>	T	F		1	1	1	1			1			5
Sapindaceae	<i>Majidea fosteri</i>	T	R		1									1
Sapindaceae	<i>Paullinia pinnata</i>	C	F		1	1		1						3
Sapindaceae	<i>Zahna golungensis</i>	C	F	1		1	1	1						4
Sapotaceae	<i>Aningeria altissima</i>	T	F	1		1			1	1	1			5
Sapotaceae	<i>Chrysophyllum albidum</i>	T	A			1	1	1	1	1	1			6
Sapotaceae	<i>Chrysophyllum muerense</i>	T	F				1	1	1		1			4
Sapotaceae	<i>Manilkara dawei</i>	T	F						1	1	1			3
Sapotaceae	<i>Mimusops bagshawei</i>	T	R			1								1
Sapotaceae	<i>Pachystela brevipes</i>	T	O					1	1					2
Simaroubaceae	<i>Harrisonia abyssinica</i>	T	R						1					1
Smilacaceae	<i>Smilax anceps</i>	C	O			1						1		2
Solanaceae	<i>Capsicum frutescens</i>	S	O	1									1	2
Solanaceae	<i>Physalis peruviana</i>	S	F			1	1	1	1					4
Solanaceae	<i>Solanum indicum</i>	S	R	1										1
Solanaceae	<i>Solanum mauritanum</i>	S	A			1	1	1	1		1	1		6
Solanaceae	<i>Sorghum arundinaceum</i>	Gr	R		1									1
Sterculiaceae	<i>Byttneria catalpifolia</i>	C	R							1				1
Sterculiaceae	<i>Cola gigantea</i>	T	O			1			1					2
Sterculiaceae	<i>Dombeya dawei</i>	T	R					1						1
Sterculiaceae	<i>Dombeya mukole</i>	T	R		1									1
Sterculiaceae	<i>Leptonychia mildbraedii</i>	T	F			1		1	1	1	1			5
Thelypteridaceae	<i>Christella parasitica</i>	H	R									1		1
Thelypteridaceae	<i>Thelypteris hamulosa</i>	H	O					1		1				2
Thymelaeaceae	<i>Peddiea fischeri</i>	H	R									1		1
Tiliaceae	<i>Grewia bicolor</i>	T	R	1										1
Tiliaceae	<i>Grewia mildbraedii</i>	T	F	1	1	1				1				4
Tiliaceae	<i>Triumfetta rhomboidea</i>	H	R	1										1
Ulmaceae	<i>Celtis africana</i>	T	A		1	1	1	1	1	1	1	1		8
Ulmaceae	<i>Celtis durandii</i>	T	A	1	1	1	1	1		1	1			7
Ulmaceae	<i>Celtis mildbraedii</i>	T	A		1	1	1	1	1	1	1	1		8

Appendix P1 (Continued)

Family	Species	Life form	Abundance	Mabira F.R. Sites								Kifu F.R.	Namyoya F.R.	Totals
				1	2	3	4	5	6	7	8	9	10	
Ulmaceae	<i>Celtis phillipensis</i>	T	F			1	1		1	1	1			5
Ulmaceae	<i>Celtis wightii</i>	T	F			1		1	1					3
Ulmaceae	<i>Celtis zenkeri</i>	T	F				1	1	1	1	1			5
Ulmaceae	<i>Holoptelea grandis</i>	T	F					1	1	1	1			4
Ulmaceae	<i>Trema orientalis</i>	T	F		1			1		1	1			4
Umbelliferae	<i>Centella asiatica</i>	H	A	1		1	1	1	1	1	1		1	8
Urticaceae	<i>Boehmeria macrophylla</i>	S	O							1	1			2
Verbenaceae	<i>Clerodendrum myricoides</i>	S	R	1										1
Verbenaceae	<i>Clerodendrum silvaticum</i>	C	R									1		1
Verbenaceae	<i>Lantana camara</i>	S	A	1	1		1	1	1	1	1		1	8
Verbenaceae	<i>Lantana triphylla</i>	S	F					1	1		1			3
Verbenaceae	<i>Lippia grandifolia</i>	S	R			1								1
Verbenaceae	<i>Stachytarpheta jamaicensis</i>	S	R			1								1
Verbenaceae	<i>Vitex amboniensis</i>	T	R				1							1
Violaceae	<i>Rinorea ilicifolia</i>	T	O				1	1						2
Vitaceae	<i>Cissus petiolata</i>	C	R	1										1
Vitaceae	<i>Cissus rotundifolia</i>	C	R		1									1
Zingiberaceae	<i>Aframomum angustifolia</i>	H	R							1				1
Zingiberaceae	<i>Aframomum mildbraedii</i>	H	R				1							1
Zingiberaceae	<i>Renealmia congolana</i>	H	O								1	1		2
Zingiberaceae	<i>Renealmia engleri</i>	H	R							1				1
Total				63	66	96	81	76	84	74	65	85	41	

Appendix P2: Plant species list for Lubigi swamp

1 means presence

H = Herb; C = Climber/Creeper; T = Tree, S = Shrub

Abundance: D – Dominant, A – Abundant, F – Frequent, O – Occasional, R - Rare

Family	Species	Life form	Abundance	Lubigi swamp sites						Totals
				1	2	3	4	5	6	
Acanthaceae	<i>Acanthus pubescens</i>	S	A	1	1	1	1	1		5
Acanthaceae	<i>Dyschoriste radicans</i>	H	R						1	1
Acanthaceae	<i>Hypoestes aristata</i>	H	R		1					1
Acanthaceae	<i>Justicia heterocarpa</i>	H	R		1					1
Anacardiaceae	<i>Pseudospondias microcarpa</i>	T	R				1			1
Apocynaceae	<i>Tabernamontana odoratissima</i>	T	R	1						1
Asclepiadaceae	<i>Cynanchum polyanthum</i>	C	O	1				1		2
Asclepiadaceae	<i>Cynanchum strigosum</i>	C	R	1						1
Asclepiadaceae	<i>Dragea sp.</i>	C	R		1					1
Asclepiadaceae	<i>Gomphocarpus fruticosus</i>	S	R	1						1
Asteraceae	<i>Ageratum conyzoides</i>	H	F	1	1	1	1			4
Asteraceae	<i>Bidens pilosa</i>	H	R						1	1
Asteraceae	<i>Conyza floribunda</i>	H	R	1						1
Asteraceae	<i>Crassocephalum sp.</i>	H	F	1	1	1		1		4
Asteraceae	<i>Enhydra fluctuans</i>	H	R						1	1
Asteraceae	<i>Melanthera scandens</i>	H	F		1	1	1	1		4
Asteraceae	<i>Mikania cordata</i>	H	R						1	1
Asteraceae	<i>Siegesbeckia abyssinica</i>	H	R						1	1
Asteraceae	<i>Spilanthes mauritiana</i>	H	R		1					1
Asteraceae	<i>Vernonia amygdalina</i>	S	F		1		1	1	1	4
Asteraceae	<i>Vernonia auriculifera</i>	H	R					1		1
Asteraceae	<i>Vernonia lasciopus</i>	H	R					1		1
Bignoniaceae	<i>Spathodea campanulata</i>	T	R	1						1
Commelinaceae	<i>Aneilema beniniense</i>	H	R			1				1
Commelinaceae	<i>Commelina africana</i>	H	O	1	1					2
Commelinaceae	<i>Commelina erecta</i>	H	O					1	1	2
Commelinaceae	<i>Commelina thomasi</i>	H	R						1	1
Convolvulaceae	<i>Ipomoea cairica</i>	C	D	1	1	1	1	1	1	6
Cyperaceae	<i>Cyperus cyperoides</i>	H	R			1				1
Cyperaceae	<i>Cyperus denudatus</i>	H	A	1		1	1	1	1	5
Cyperaceae	<i>Cyperus distans</i>	H	R	1						1
Cyperaceae	<i>Cyperus latifolius</i>	H	A	1	1	1	1		1	5
Cyperaceae	<i>Cyperus papyrus</i>	H	A	1	1	1	1		1	5
Cyperaceae	<i>Cyperus sp.</i>	H	R		1					1
Cyperaceae	<i>Fimbristylis dichotoma</i>	H	F	1		1	1			3
Cyperaceae	<i>Fimbristylis miliaceae</i>	H	O				1	1		2
Cyperaceae	<i>Kyllinga sp.</i>	H	D	1	1	1	1	1	1	6
Cyperaceae	<i>Pycnus flavescens</i>	H	R	1						1
Cyperaceae	<i>Pycnus nitidus</i>	H	D	1	1	1	1	1	1	6
Cyperaceae	<i>Scleria achenii</i>	H	A	1	1	1	1	1		5
Cyperaceae	<i>Scleria bulbifera</i>	H	R			1				1
Cyperaceae	<i>Scleria catophylla</i>	H	O			1	1			2
Cyperaceae	<i>Scleria melanomphala</i>	H	A	1	1	1	1	1		5
Cyperaceae	<i>Scleria niasensis</i>	H	O	1	1					2
Davalliaceae	<i>Nephrolepis biserrata</i>	H	R	1						1
Euphorbiaceae	<i>Alchornea cordifolia</i>	T	O	1		1				2
Euphorbiaceae	<i>Bridelia micrantha</i>	T	F		1		1		1	3
Euphorbiaceae	<i>Euphorbia sp.</i>	H	O			1			1	2

Appendix P2 (Continued)

Family	Species	Life form	Abundance	Lubigi swamp sites						Totals
				1	2	3	4	5	6	
Euphorbiaceae	<i>Phyllanthus nummulariifolius</i>	H	F			1		1	1	3
Euphorbiaceae	<i>Phyllanthus ovalifolius</i>	H	R			1				1
Euphorbiaceae	<i>Ricinus communis</i>	S	O					1	1	2
Fabaceae	<i>Aeschynomene indica</i>	H	F			1	1	1	1	4
Fabaceae	<i>Aeschynomene schimperi</i>	S	R		1					1
Fabaceae	<i>Albizia grandibracteata</i>	T	F			1	1	1		3
Fabaceae	<i>Albizia zygia</i>	T	R	1						1
Fabaceae	<i>Cassia kirki</i>	H	O			1		1		2
Fabaceae	<i>Cassia mimosoides</i>	H	O	1				1		2
Fabaceae	<i>Crotalaria cleomifolia</i>	H	R				1			1
Fabaceae	<i>Crotalaria ochroleuca</i>	H	R	1						1
Fabaceae	<i>Desmodium ramosissimum</i>	H	R	1						1
Fabaceae	<i>Desmodium salicifolium</i>	H	F	1	1	1			1	4
Fabaceae	<i>Desmodium velutinum</i>	H	O					1	1	2
Fabaceae	<i>Eriosema laurentii</i>	H	O			1	1			2
Fabaceae	<i>Erythrina abyssinica</i>	T	F			1	1	1		3
Fabaceae	<i>Indigofera spicata</i>	H	O					1	1	2
Fabaceae	<i>Kotschya africana</i>	H	O					1	1	2
Fabaceae	<i>Mimosa pigra</i>	H	A	1		1	1	1	1	5
Fabaceae	<i>Vigna luteola</i>	H	A	1	1		1	1	1	5
Fabaceae	<i>Vigna parkeri</i>	H	R	1						1
Fabaceae	<i>Vigna sp.</i>	H	R	1						1
Guttiferae	<i>Harungana madagascariensis</i>	T	F			1	1	1		3
Labiatae	<i>Geniosporum rotundifolium</i>	H	R				1			1
Labiatae	<i>Hyptis lanceolata</i>	H	F	1			1		1	3
Labiatae	<i>Leonotis nepetifolia</i>	H	F			1	1	1		3
Labiatae	<i>Ocimum grattisimum</i>	H	R				1			1
Labiatae	<i>Plectranthus sp.</i>	H	O			1	1			2
Malvaceae	<i>Hibiscus diversifolius</i>	H	A	1	1		1	1	1	5
Malvaceae	<i>Sida rhombifolia</i>	H	O					1	1	2
Malvaceae	<i>Sida sp.</i>	H	F		1		1	1		3
Melastomataceae	<i>Dissotis canescens</i>	H	R					1		1
Melastomataceae	<i>Dissotis rotundifolia</i>	H	R		1					1
Melastomataceae	<i>Tristemma mauritianum</i>	H	F	1	1				1	3
Menispermaceae	<i>Cissampelos mucronata</i>	C	D	1	1	1	1	1	1	6
Menispermaceae	<i>Stephania abyssinica</i>	H	R						1	1
Moraceae	<i>Antiaris toxicaria</i>	T	R	1						1
Moraceae	<i>Ficus ovata</i>	T	R	1						1
Moraceae	<i>Ficus vallis-chaude</i>	T	O	1		1				2
Moraceae	<i>Ficus verruculosa</i>	T	R	1						1
Myrsinaceae	<i>Maesa lanceolata</i>	S	R	1						1
Myrtaceae	<i>Eucalyptus grandis</i>	T	F	1	1				1	3
Myrtaceae	<i>Syzygium guineense</i>	T	R			1				1
Onagraceae	<i>Jussiaea abyssinica</i>	H	R	1						1
Orchidaceae	<i>Eulophia horsfallii</i>	H	R			1				1
Palmae	<i>Phoenix reclinata</i>	T	A	1	1	1	1	1		5
Poaceae	<i>Brachiaria decumbens</i>	G	R			1				1
Poaceae	<i>Chloris sp.</i>	G	O			1			1	2
Poaceae	<i>Cymbopogon sp.</i>	G	R			1				1
Poaceae	<i>Digitaria abyssinica</i>	G	F			1	1	1	1	4
Poaceae	<i>Eragrostis mildbraedii</i>	G	R	1						1

Appendix P2 (Continued)

Family	Species	Life form	Abundance	Lubigi swamp sites						Totals
				1	2	3	4	5	6	
Poaceae	<i>Hyparrhenia sp.</i>	G	O			1		1		2
Poaceae	<i>Leersia hexandra</i>	G	D	1	1	1	1	1	1	6
Poaceae	<i>Loudetia kagerensis</i>	G	R					1		1
Poaceae	<i>Loudetia phragmatoides</i>	G	R					1		1
Poaceae	<i>Miscanthus violaceus</i>	G	R					1		1
Poaceae	<i>Panicum maximum</i>	G	A	1	1	1		1	1	5
Poaceae	<i>Setaria sphacelata</i>	H	F	1		1	1	1		4
Poaceae	<i>Sporobolus sp.</i>	G	R						1	1
Polygonaceae	<i>Polygonum salicifolium</i>	H	F		1		1	1	1	4
Polygonaceae	<i>Polygonum sp.</i>	H	O	1	1					2
Polygonaceae	<i>Polygonum strigosum</i>	H	R	1						1
Primulaceae	<i>Lysimachia ruhmeriana</i>	H	R	1						1
Rosaceae	<i>Rubus apetalus</i>	S	F	1		1	1			3
Sapindaceae	<i>Paullinia pinnata</i>	C	R	1						1
Schizaeaceae	<i>Lygodium microphyllum</i>	H	R		1					1
Smilacaceae	<i>Smilax anceps</i>	C	R			1				1
Solanaceae	<i>Solanum mauritianum</i>	S	O			1	1			2
Thelypteridaceae	<i>Thelypteris confluens</i>	H	O					1	1	2
Thelypteridaceae	<i>Thelypteris fadenii</i>	H	R	1						1
Thelypteridaceae	<i>Thelypteris totta</i>	H	A	1	1	1	1		1	5
Tiliaceae	<i>Triumfetta macrophylla</i>	S	A		1	1	1	1	1	5
Typhaceae	<i>Typha domingensis</i>	H	F	1		1	1		1	4
Verbenaceae	<i>Clerodendrum fuseum</i>	C	R				1			1
Verbenaceae	<i>Lantana camara</i>	S	F			1	1	1		3
Zingiberaceae	<i>Aframomum angustifolium</i>	H	F	1		1	1	1		4
Totals				57	37	51	45	47	42	

Appendix A1 Dragonfly species list

Species	Forest sites										Swamp sites							
	Mabira																	
										Kifu	Namyoya							
	8	7	6	5	4	3	2	1				1	2	3	4	5	6	Typical Habitat
Zygoptera																		
<i>Chlorophora trifaria</i>						1												thickly forested streams
<i>Platycypha lacustris</i>		1		1								1						rivers and streams in dense forest
<i>Umma saphirina</i>						1		1	1									thickly forested streams
Anisoptera																		
<i>Gynacantha villosa</i>				1												1	1	forest, thick bush
<i>Orthetrum julia</i>		1		1		1												forest, dense woodland, streams
<i>Orthetrum macrostigma</i>			1											1	1		1	bush fringed swamps and pools
<i>Orthetrum microstigma</i>			1															swamp forest
<i>Orthetrum</i> sp 1			1															
<i>Orthetrum</i> sp 2						1							1		1		1	
<i>Orthetrum</i> sp 3										1			1					
<i>Orthetrum</i> sp 4								1							1		1	
<i>Orthetrum</i> sp 5			1															
<i>Orthetrum trinacria</i>										1			1		1	1	1	pools, lakes, rivers in savannah, bush, woodland
<i>Palpopleura lucia</i>		1						1										reedy sluggish streams and pools in woodland and forest
<i>Palpopleura portia</i>		1						1										reedy or grassy sluggish streams or pools in savannah, woodland, bush
<i>Pseudagrion kersteni</i>						1						1	1	1	1	1	1	abundant in most habitat except dense forest
<i>Pseudagrion melanicterum</i>						1												forest streams, thich bush and litmus
<i>Pseudagrion rufocinctum</i>			1			1												forest
<i>Pseudagrion</i> sp						1							1			1		
<i>Pseudagrion spermatum</i>			1															montane streams or rivers, shade or thick bush
Totals	0	4	6	3	0	8	0	0	4	3	2	5	2	5	4	6		
All forest and swamp sites						20								9				

Appendix A2. Butterfly species list

Species	Forest sites										Swamp sites						Ecotype ^a
	8	7	6	5	4	3	2	1	Kifu	Namyoya	1	2	3	4	5	6	
Nymphalidae																	
<i>Acraea bonasia</i>											1	1					F
<i>Acraea acerata</i>									1			1					W
<i>Acraea aganice</i>	1	1	1														f
<i>Acraea alicia</i>											1	1					W
<i>Acraea althoffi</i>	1								1								F
<i>Acraea aurivillii</i>	1				1	1			1								F
<i>Acraea cerasa</i>									1								f
<i>Acraea egina</i>		1	1	1		1											W
<i>Acraea encedon</i>							1	1	1		1	1					W
<i>Acraea epaea</i>	1																F
<i>Acraea eponina</i>			1	1	1	1	1	1	1					1			W
<i>Acraea humilis</i>	1																F
<i>Acraea jodutta</i>	1				1												F
<i>Acraea johnstoni</i>	1				1			1						1			f
<i>Acraea leucographa</i>							1							1			F
<i>Acraea lycoa</i>		1		1	1	1			1								F
<i>Acraea macaria</i>	1						1										F
<i>Acraea macarista</i>								1						1			F
<i>Acraea natalica</i>		1		1			1										W
<i>Acraea neobule</i>				1						1							W
<i>Acraea orinata</i>					1	1											F
<i>Acraea peneleos</i>						1		1									F
<i>Acraea penelope</i>		1			1	1											F
<i>Acraea poggei</i>									1								F
<i>Acraea psudegina</i>		1	1	1		1	1			1	1	1	1	1	1	1	W
<i>Acraea quirina</i>								1	1								F
<i>Acraea quirinalis</i>	1			1					1								F
<i>Acraea rogersi</i>	1	1	1	1													F
<i>Acraea servona</i>	1																F
<i>Acraea tellus</i>								1									F
<i>Acraea uvui</i>									1				1	1			f
<i>Acraea venrura</i>																	S
<i>Acraea viviana</i>	1			1					1	1							f
<i>Acraea zetes</i>		1							1								W
<i>Amauris albimaculata</i>	1						1			1							F
<i>Amauris echeria</i>							1										f
<i>Amauris niavius</i>		1							1	1							W
<i>Amauris oscarus</i>				1										1			F
<i>Tirumala petiverana</i>						1								1			W/m

Appendix A2 (Continued)

Species	Forest sites										Swamp sites						Ecotype ^a
	8	7	6	5	4	3	2	1	Kifu	Namyoya	1	2	3	4	5	6	
<i>Amauris tartarea</i>	1	1	1				1		1	1	1	1	1	1	1	1	F
<i>Ariandne albifascia</i>	1			1			1	1	1								F
<i>Ariandne enotrea</i>								1	1	1							f
<i>Aterica galene</i>							1		1								F
<i>Bebearia ribensis</i>									1								F
<i>Bicyclus auricrudus</i>	1	1						1	1								F
<i>Bicyclus funebris</i>	1								1								FL
<i>Bicyclus istaris</i>									1					1	1		f
<i>Bicyclus jefferyi</i>		1			1	1			1	1	1	1	1	1	1	1	f
<i>Bicyclus mesogena</i>	1	1															F
<i>Bicyclus mollitia</i>	1	1															F
<i>Bicyclus sambulos</i>	1	1															F
<i>Bicyclus sandace</i>		1		1	1	1	1	1									F
<i>Bicyclus saussurei</i>																	F
<i>Bicyclus sebetus</i>	1				1												FL
<i>Bicyclus safitza</i>										1							W
<i>Bicyclus smithi</i>		1			1			1	1								FL
<i>Bicyclus sophrosyne</i>				1	1		1	1									f
<i>Bicyclus uniformis</i>	1	1			1												FL
<i>Bicyclus vulgaris</i>					1	1		1	1		1						W
<i>Byblia anvatarata</i>	1	1		1	1	1	1		1	1	1	1	1	1	1	1	W/m
<i>Byblia ilithyia</i>									1								O
<i>Catuna crithea</i>	1	1			1	1	1										F
<i>Charaxes acuminatus</i>		1															FH
<i>Charaxes cynthia</i>	1				1												F
<i>Charaxes etheocles</i>						1											F
<i>Charaxes fulvescence</i>					1				1								FL
<i>Charaxes pleione</i>						1											f
<i>Charaxes tiridates</i>						1											FL
<i>Charaxes varanes</i>					1	1											W
<i>Charaxes zelica</i>						1											F
<i>Cyrestis camillus</i>						1											F
<i>Danaus chrysippus</i>						1	1		1	1	1						O/m
<i>Euphaedra eleus</i>	1	1			1	1											F
<i>Euphaedra harpalyce</i>					1												F
<i>Euphaedra medon</i>	1				1	1			1								F
<i>Euphaedra preussi</i>	1	1															F
<i>Euphaedra rex</i>	1	1															F
<i>Euphaedra ruspina</i>						1	1										F

Appendix A2 (Continued)

Species	Forest sites										Swamp sites						Ecotype ^a
	8	7	6	5	4	3	2	1	Kifu	Namyoya	1	2	3	4	5	6	
<i>Eurytela dryope</i>									1	1							W
<i>Eurytela hiarbas</i>			1		1	1	1	1	1								f
<i>Gnophodes betsimena</i>		1															F
<i>Gnophodes chelys</i>		1			1		1										F
<i>Harma theobene</i>	1			1		1		1									F
<i>Henotesia peitho</i>		1			1												W
<i>Henotesia perspicua</i>											1						O
<i>Hypolimnias monteironis</i>	1																F
<i>Hypolimnias salmacis</i>	1						1										F
<i>Junonia chorimene</i>								1		1							O
<i>Junonia oenone</i>							1										W
<i>Junonia sophia</i>				1	1	1	1	1	1	1	1	1	1	1	1	1	W
<i>Junonia stygia</i>						1											f
<i>Junonia terea</i>				1				1		1							W
<i>Junonia westermanni</i>	1	1		1	1	1	1		1								F
<i>Libythea labdaca</i>	1			1		1											W/m
<i>Melanitis leda</i>						1											W
<i>Mesoxantha ethosea</i>	1	1		1													FL
<i>Neptidopsis ophione</i>				1				1	1								f
<i>Neptis melicerta</i>			1			1		1									F
<i>Neptis metella</i>				1			1							1			f
<i>Neptis necomedes</i>						1											f
<i>Neptis nemetes</i>					1	1			1								f
<i>Neptis ochracea</i>									1								F
<i>Neptis trigonophora</i>							1										F
<i>Neptis saclava</i>				1		1	1	1	1								W
<i>Neptis serena</i>					1						1						W
<i>Phalanta phalanta</i>	1							1					1				O/m
<i>Pseudacraea deludens</i>							1										FH
<i>Pseudacraea lucretia</i>				1	1	1											f
<i>Pseudoneptis bugandensis</i>	1					1											F
<i>Salamis cacta</i>							1										F
<i>Salamis parhassus</i>				1		1			1								f
<i>Sallya boisduvali</i>						1											f/m

Appendix A2 (Continued)

Species	Forest sites								Swamp sites								Ecotype ^a
	8	7	6	5	4	3	2	1	Kifu	Namyoya	1	2	3	4	5	6	
<i>Sallya garega</i>				1													F/m
<i>Sallya occidentarium</i>						1											F/m
<i>Tirumala petiverana</i>		1															W/m
<i>Venessula milca</i>						1											f
<i>Ypthima albida</i>					1	1			1								f
<i>Ypthima antennata</i>											1				1		O
<i>Ypthima asterope</i>						1	1	1					1				O
<i>Ypthima doleta</i>									1				1			1	W
<i>Ypthimomorpha itonia</i>											1						f
Pieridae																	
<i>Appias epaphia</i>	1							1			1						f/m
<i>Appias sabina</i>							1	1									F
<i>Appias sylvia</i>						1											F
<i>Belenois aurota</i>													1	1			O/m
<i>Belenois calypso</i>	1			1		1											F
<i>Belenois crawshayi</i>							1	1									F
<i>Belenois creona</i>							1			1							O/m
<i>Belenois solilucis</i>						1											f
<i>Belenois theora</i>				1	1	1											f
<i>Belenois thysa</i>				1				1	1								f
<i>Catopsilia florella</i>				1	1												O/m
<i>Dixeia orbona</i>	1			1			1	1									W
<i>Eurema desjardinsi</i>											1				1		W
<i>Eurema hapale</i>											1		1				S
<i>Eurema hecabe</i>	1	1			1			1	1	1							W/m
<i>Eurema senegalensis</i>							1		1								F
<i>Leptosia alcesta</i>					1						1						W
<i>Leptosia hybrida</i>	1	1		1	1		1										F
<i>Leptosia nupta</i>	1			1	1	1			1								F
<i>Leptosia wigginsi</i>	1		1	1	1	1	1	1									F
<i>Mylothris continua</i>					1	1											F
<i>Mylothris hilara</i>						1											F
<i>Mylothris rubricosta</i>											1				1		S
<i>Mylothris schumanni</i>						1											F
<i>Nepheronia argia</i>	1			1	1		1		1								F
<i>Nepheronia pharis</i>							1										F

Appendix A2 (Continued)

Species	Forest sites										Swamp sites						Ecotype ^a
	8	7	6	5	4	3	2	1	Kifu	Namyoya	1	2	3	4	5	6	
<i>Nepheronia thalassina</i>	1			1	1	1											F
Lycaenidae																	
<i>Abisara neavei</i>	1			1	1	1		1	1								F
<i>Anthene indefinita</i>					1	1											O
<i>Anthene larydas</i>						1											F
<i>Anthene princeps</i>					1	1											O
<i>Anthene schoutedeni</i>				1	1				1								F
<i>Cupidopsis jobates</i>											1						W
<i>Eicochrysops hippocrates</i>														1			W
<i>Eicochrysops messapus</i>													1				O
<i>Epitola mpanensis</i>									1								F
<i>Euchrysops malathana</i>											1			1			O
<i>Hypolycaena liara</i>	1																F
<i>Hypolycaena philippus</i>						1			1								W
<i>Iolaus parasilanus</i>						1											F
<i>Larinopoda tera</i>									1								F
<i>Leptotes pirithous</i>						1	1										W/m
<i>Liptena xanthostola</i>				1													F
<i>Megalopalpus zymna</i>				1	1	1			1								F
<i>Mimeresia sp</i>									1								F
<i>Oboronia punctatus</i>						1			1								F
<i>Phlyaria heritsia</i>													1				F
<i>Tetrarhanis ilma</i>					1		1		1								F
<i>Thermoniphas micyclus</i>											1			1			F
<i>Triclema nigeriae</i>									1								f
<i>Tuxentius margaritaceus</i>						1			1								W
<i>Uranothauma falkensteini</i>		1				1											W
<i>Zizeeria knysna</i>							1				1						1W
<i>Zizina antanossa</i>											1						W
<i>Zizula hylax</i>							1										W
Papilionidae																	
<i>Papilio bromius</i>	1			1	1	1	1		1								f
<i>Papilio cynorta</i>	1				1	1											FL
<i>Papilio dardanus</i>	1		1	1	1	1		1	1								W

Appendix A2 (Continued)

Species	Forest sites										Swamp sites						Ecotype ^a
	8	7	6	5	4	3	2	1	Kifu	Namyoya	1	2	3	4	5	6	
<i>Papilio demodocus</i>				1	1		1	1	1	1							W/m
<i>Papilio echerioides</i>									1								f
<i>Papilio interjecta</i>					1												F
<i>Papilio lormieri</i>	1				1												F
<i>Papilio nireus</i>					1												f
<i>Papilio nobilis</i>				1													F
<i>Papilio phorcas</i>	1	1	1	1	1	1	1	1									F
<i>Papilio rex</i>		1			1												FH
Hesperiidae																	
<i>Acleros ploetzi</i>																	f
<i>Andronymus neander</i>			1														f/m
<i>Ankola fan</i>											1			1		1	F
<i>Borbo fallax</i>						1			1								O
<i>Borbo kaka</i>														1			F
<i>Borbo lugens</i>									1		1						f
<i>Borbo micans</i>											1			1			S
<i>Calaenorrhinus proxima</i>	1																F
<i>Calaenorrhinus bettoni</i>					1												f
<i>Calaenorrhinus galenus</i>	1					1		1									F
<i>Calaenorrhinus intermixtus</i>									1								F
<i>Ceratrachia flava</i>						1											F
<i>Coeliades forestan</i>							1	1	1	1							W
<i>Eagris lucetia</i>									1								f
<i>Eretis lugens</i>	1						1		1	1							W
<i>Gegenes hottentota</i>														1			O
<i>Gegenes niso</i>	1					1	1										W
<i>Gorgyra sp</i>											1						f
<i>Lepella lepeletier</i>														1			f
<i>Metisella midas</i>											1			1		1	S
<i>Monza alberti</i>	1																F
<i>Pardeleodes incerta</i>				1						1					1		F
<i>Pardeleodes tibullus</i>					1	1	1										F
<i>Sarangesa bouvieri</i>	1																F
<i>Spialia spio</i>	1										1						O

Note a : see Table A3

Appendix A3. Comparison of butterfly species recorded from Mabira forest.

Species	Previous record only	This study but not previously
Nymphalidae		
<i>A. homilis</i>		X
<i>A. neobule</i>		X
<i>A. pharsalus</i>	X	
<i>A. quirina</i>		X
<i>A. zetes</i>		X
<i>A.leucographa</i>		X
<i>A.viviana</i>		X
<i>Acraea cabira</i>	X	
<i>Acraea encedon</i>		X
<i>Amauris echeria</i>		X
<i>Amauris oscarus</i>		X
<i>Antanartia delius</i>	X	
<i>Ariadne pagenstecheri</i>	X	
<i>Bebearia cocalia</i>	X	
<i>Bicyclus campinus</i>	X	
<i>Bicyclus sebetus</i>	X	
<i>Charaxes ameliae</i>	X	
<i>Charaxes bipunctatus</i>	X	
<i>Charaxes brutus</i>	X	
<i>Charaxes candiope</i>	X	
<i>Charaxes etesipe</i>	X	
<i>Charaxes eupale</i>	X	
<i>Charaxes lucretius</i>	X	
<i>Charaxes numenes</i>	X	
<i>Charaxes porthos</i>	X	
<i>Charaxes protoclea</i>	X	
<i>Charaxes pythodorus</i>	X	
<i>Charaxes subornatus</i>	X	
<i>Charaxes virilis</i>	X	
<i>Charaxes zelica</i>		X
<i>Charaxes zingha</i>	X	
<i>Cymothoe caenis</i>	X	
<i>Cymothoe herminia</i>	X	
<i>Cymothoe hobarti</i>	X	
<i>Euphaedra ruspina</i>		X
<i>Euphaedra uganda</i>	X	
<i>Euryphura albimargo</i>	X	
<i>Euryphura chalcis</i>	X	
<i>Euxanthe crossleyi</i>	X	
<i>Gnophodes chelys</i>		X
<i>Hypolimnas dinarcha</i>	X	
<i>Hypolimnas dubius</i>	X	
<i>Hypolimnas misippus</i>	X	
<i>Hypolimnas monteironis</i>		X
<i>Junonia oenone</i>		X
<i>Neptis conspicua</i>	X	
<i>Neptis serena</i>		X
<i>Palla ussheri</i>	X	
<i>Phalanta eurytis</i>	X	
<i>Phalanta phalanta</i>		X

Appendix A3 (Continued)

Species	Previous record only	This study but not previously
<i>Pseudacraea clarki</i>	x	
<i>Pseudacraea deludens</i>		x
<i>Pseudacraea eurytus</i>	x	
<i>Pseudathyma plutonica</i>	x	
<i>Salamis cacta</i>	x	
<i>Sallya natalensis</i>	x	
<i>Tirumala formosa</i>	x	
<i>Venessula milca</i>		x
<i>Ariadne albifascia</i>		x
Pieridae		
<i>Belenois aurota</i>		x
<i>Belenois crawshayi</i>		x
<i>Catopsilia florella</i>		x
<i>Dixeia charina</i>	x	
<i>Eurema desjardinsi</i>		x
<i>Colotis elgonensis</i>	x	
<i>Belenois subeida</i>	x	
<i>Mylothris rubricosta</i>		x
<i>Mylothris hilara</i>		x
<i>Mylothris schumanni</i>		x
Lycaenidae		
<i>Abisara neavei</i>		x
<i>Anthene definita</i>	x	
<i>Anthene lingures</i>	x	
<i>Anthene princeps</i>		x
<i>Azanus isis</i>	x	
<i>Azanus jesous</i>	x	
<i>Azanus mirza</i>	x	
<i>Azanus moriqua</i>	x	
<i>Azanus natalensis</i>	x	
<i>Cacyreus audeoudi</i>	x	
<i>Cacyreus lingeus</i>	x	
<i>Epitola catuna</i>	x	
<i>Hypolycaena antifaunus</i>	x	
<i>Hypolycaena hatita</i>	x	
<i>Hypolycaena liara</i>		x
<i>Iolaus parasılanus</i>		x
<i>Lachnocnema bibulus</i>	x	
<i>Leptotes pirithous</i>		x
<i>Liptena xanthostola</i>		x
<i>Pentila pauli</i>	x	
<i>Phylaria cyara</i>	x	
<i>Phylaria heritsia</i>		x
<i>Tanuetheira timon</i>	x	
<i>Tetrarhanis ilma</i>		x
<i>Tuxentius cretosus</i>	x	
<i>Zizeeria knysna</i>		x
<i>Zizina antanossa</i>		x
<i>Zizula hylax</i>		x
Papilionidae		
<i>Graphium polices</i>	x	
<i>Papilio echerioides</i>	x	

Appendix A3 (Continued)

Species	Previous record only	This study but not previously
<i>Papilio interjescta</i>		X
<i>Papilio nobilis</i>		X
<i>Papilio rex</i>		X
Hesperiidae		
<i>Acleros mackenii</i>	X	
<i>Andronymus neander</i>		X
<i>Borbo fallax</i>		X
<i>Borbo gemella</i>	X	
<i>Caenides dacena</i>	X	
<i>Celaenorrhinus bettoni</i>		X
<i>Ceratrachia flava</i>		X
<i>Ceratrachia mabirensis</i>	X	
<i>Gegenes niso</i>		X
<i>Gomalia elma</i>	X	
<i>Monza alberti</i>		X
<i>Monza cretacea</i>	X	
<i>Pardeleodes incerta</i>		X
<i>Pardeleodes tibullus</i>		X
<i>Sarangesa lucidella</i>	X	
<i>Spialia spio</i>		X

Appendix B1. Summary of bird count data from Mabira forest. The columns T1-8 and M1-8 are the TSC scores, and numbers of birds mist-netted, respectively. Mean values are also given.

The column OP lists a few additional species that were recorded opportunistically

Atlas No ^a	Species	Status ^b	Spec ^c	RD ^d	T1	T1	T2	T3	T4	T5	T6	T7	T8	Mean TSC	M1	M2	M3	M4	M5	M6	M7	M8	Mean Net	OP
26	BLACK-HEADED HERON <i>Ardea melanocephala</i>	RB	w		0	0	0	0	0	0	0	0	0		0	0	0	0	0	0	0		1	
86	BROWN SNAKE-EAGLE <i>Circaetus cinereus</i>	R(B)		R-NT	0	0	0	0	0	0	0	0	0		0	0	0	0	0	0	0		1	
109	LIZARD BUZZARD <i>Kaupifalco monogrammicus</i>	RB	F		0	0	0	0	0	0	0	6	0	0.9	0	0	0	0	0	0	0	0.0	0	
124	CROWNED EAGLE <i>Stephanoaetus coronatus</i>	RB	FF	R-VU	0	0	0	0	0	0	0	0	0		0	0	0	0	0	0	0		1	
141	CRESTED GUINEAFOWL <i>Guttera pucherani</i>	RB	F		0	0	6	0	0	0	0	0	0	1.2	0	0	0	0	0	0	0	0.0	0	
155	SCALY FRANCOLIN <i>Francolinus squamatus</i>	RB	F		0	0	0	0	0	0	0	0	0		0	0	0	0	0	0	0		1	
156	NAHAN'S FRANCOLIN <i>Francolinus nahani</i>	RB	FF	G-ED	0	0	0	0	0	5	0	0	0	0.5	0	0	0	0	0	0	0	0.0	0	
168	WHITE-SPOTTED FLUEFTAIL <i>Sarothrura pulchra</i>	RB	FW		0	0	0	5	5	0	3	0	4	2.2	0	0	0	0	0	0	0	0.0	0	
268	AFRICAN GREEN-PIGEON <i>Treron calva</i>	RB	F		0	0	0	0	0	0	0	0	0		0	0	0	0	0	0	0		1	
270	TAMBOURINE DOVE <i>Turtur tympanistria</i>	RB	F		6	6	6	6	0	6	3	0	5	5.0	0	0	0	0	0	0	1	0	0.1	0
271	BLUE-SPOTTED WOOD-DOVE <i>Turtur afer</i>	RB	F		0	0	5	0	0	5	6	0	6	2.6	0	0	1	0	0	0	0	0.1	0	
280	AFEP PIGEON <i>Columba unicincta</i>	RB	FF		0	0	0	0	0	0	3	6	1	1.0	0	0	0	0	0	0	0	0.0	0	
283	RED-EYED DOVE <i>Streptopelia semitorquata</i>	RB	f		0	0	0	0	0	0	0	0	0		0	0	0	0	0	0	0		1	
290	GREY PARROT <i>Psittacus erithacus</i>	RB	FF	R-NT	0	0	0	0	0	0	5	0	0	1.0	0	0	0	0	0	0	0	0.0	0	
296	GREAT BLUE TURACO <i>Corythaeola cristata</i>	RB	F		0	0	0	5	2	5	0	2	0	1.5	0	0	0	0	0	0	0	0.0	0	
297	BLACK-BILLED TURACO <i>Tauraco schuetti</i>	RB	FF		0	0	0	1	0	0	2	0	0	0.8	0	0	0	0	0	0	0	0.0	0	
309	RED-CHESTED CUCKOO <i>Cuculus solitarius</i>	RB, AfM/NB?	AF		0	0	0	0	0	0	5	0	1	0.6	0	0	0	0	0	0	0	0.0	0	
310	BLACK CUCKOO <i>Cuculus clamosus</i>	RB? AfM/B?	Af/FF		0	0	0	0	0	0	0	0	0		0	0	0	0	0	0	0		1	
314	DUSKY LONG-TAILED CUCKOO <i>Cercococcyx mechowi</i>	RB	FF		0	0	0	5	0	0	3	0	4	1.6	0	0	1	0	0	0	0	0.1	0	
317	AFRICAN EMERALD CUCKOO <i>Chrysococcyx cupreus</i>	RB?	F		0	0	0	1	0	1	3	0	3	1.6	0	0	0	0	0	0	0	0.0	0	
319	KLAAS' CUCKOO <i>Chrysococcyx klaas</i>	RB	f		0	0	0	0	0	0	4	0	2	1.1	0	0	0	0	0	0	0	0.0	0	
321	YELLOWBILL <i>Ceuthmochares aereus</i>	RB	F		0	0	0	0	0	0	0	6	0	0.7	0	0	0	0	0	0	0	0.0	0	
323	WHITE-BROWED COUCAL <i>Centropus superciliosus</i>	RB			0	0	0	0	0	4	0	0	0	1.0	0	0	0	0	0	0	0	0.0	0	
358	AFRICAN PALM SWIFT <i>Cypsiurus parvus</i>	RB			0	0	0	0	0	0	0	0	0	0.1	0	0	0	0	0	0	0	0.0	0	
367	ALPINE SWIFT <i>Apus melba</i>	RB	p		0	0	0	0	0	0	0	0	0	0.4	0	0	0	0	0	0	0	0.0	0	
371	NARINA'S TROGON <i>Apaloderma narina</i>	RB	F		0	0	0	0	0	1	1	5	0	1.4	0	0	0	0	0	0	0	0.0	0	
374	BLUE-BREASTED KINGFISHER <i>Halcyon malimbica</i>	RB	Fw		0	0	0	4	0	0	0	4	0	0.8	0	0	0	0	0	0	0	0.0	0	
378	AFRICAN PYGMY KINGFISHER <i>Ceyx picta</i>	RB, AfM/NB	fw		0	0	0	0	0	0	0	0	0	0.0	1	1	2	2	0	3	0	1.1	0	
390	WHITE-THROATED BEE-EATER <i>Merops albicollis</i>	AfM/NB, FB, PM	Af		3	4	5	0	1	0	0	0	0	1.1	0	0	0	3	0	0	0	0.4	0	
419	CROWNED HORNBILL <i>Tockus alboterminatus</i>	RB	f		0	0	0	0	0	0	0	0	0		0	0	0	0	0	0	0		1	
422	BLACK-AND-WHITE CASQUED HORNBILL <i>Bycanistes subcylindricus</i>	RB	F		0	0	0	1	0	6	6	2	1	2.7	0	0	0	0	0	0	0	0.0	0	

Atlas No ^a	Species	Status ^b	Spec ^c	RD ^d	T1	T1	T2	T3	T4	T5	T6	T7	T8	Mean TSC	M1	M2	M3	M4	M5	M6	M7	M8	Mean Net	OP
426	SPECKLED TINKERBIRD <i>Pogoniulus scolopaceus</i>	RB	F		0	0	6	6	5	6	6	5	5	4.5	0	0	0	0	0	0	0	0	0.0	0
430	YELLOW-THROATED TINKERBIRD <i>Pogoniulus subsulphureus</i>	RB	FF		6	5	6	6	5	6	6	3	6	5.4	0	0	0	0	0	0	0	0	0.0	0
431	YELLOW-RUMPED TINKERBIRD <i>Pogoniulus bilineatus</i>	RB	F		0	0	0	0	0	0	3	0	0	1.5	0	0	0	0	0	0	0	0	0.0	0
434	YELLOW-SPOTTED BARBET <i>Buccanodon duchaillui</i>	RB	FF		6	1	1	4	0	6	3	0	0	2.3	0	0	0	0	0	0	0	0	0.0	0
435	HAIRY-BREASTED BARBET <i>Tricholaema hirsuta</i>	RB	F		0	4	4	5	4	5	0	4	1	2.8	0	0	0	0	0	0	0	0	0.0	0
445	YELLOW-BILLED BARBET <i>Trachyphonus purpuratus</i>	RB	FF		0	4	2	4	1	2	5	5	6	3.6	0	0	0	0	0	0	0	0	0.0	0
456	LESSER HONEYGUIDE <i>Indicator minor</i>	RB	f		0	0	0	0	0	0	0	0	0	0.0	0	1	0	0	0	0	0	0	0.1	0
469	BUFF-SPOTTED WOODPECKER <i>Campethera nivos</i>	R(B)	FF		0	0	0	0	0	0	0	0	0	0.0	0	1	0	0	2	0	0	1	0.5	0
470	BROWN-EARED WOODPECKER <i>Campethera caroli</i>	RB	FF		0	0	0	0	0	0	0	0	0	0.0	0	1	0	0	0	0	0	0	0.1	0
498	WHITE-HEADED SAW-WING <i>Psalidoprocne albiceps</i>	RB, AfM/NB?	f	R-RR	4	0	0	0	0	0	0	6	1	1.4	0	0	0	0	0	0	0	0	0.0	0
538	LITTLE GREENBUL <i>Andropadus virens</i>	RB	F		6	6	6	6	6	5	3	6	6	5.5	6	7	2	4	5	1	1	4	3.8	0
540	CAMEROON SOMBRE GREENBUL <i>Andropadus curvirostris</i>	RB	FF		0	0	0	0	0	0	0	0	0	0.0	0	0	0	1	0	0	0	0	0.1	0
541	SLENDER-BILLED GREENBUL <i>Andropadus gracilirostris</i>	RB	FF		0	0	0	0	0	0	0	0	0	0.0	0	0	0	0	0	0	2	0	0.3	0
542	YELLOW-WHISKERED GREENBUL <i>Andropadus latirostris</i>	RB	F		5	0	3	5	5	0	6	5	4	3.4	4	3	1	3	8	3	3	3	3.5	0
543	HONEYGUIDE GREENBUL <i>Baeopogon indicator</i>	R(B)	FF		0	0	0	0	0	0	0	0	0		0	0	0	0	0	0	0		1	
551	TORO OLIVE GREENBUL <i>Phyllastrephus hypochloris</i>	RB	FF	R-VU/RR	0	0	0	0	0	0	0	0	0	0.0	0	0	1	2	0	0	0	2	0.6	0
556	WHITE-THROATED GREENBUL <i>Phyllastrephus albigularis</i>	RB	FF		0	0	0	0	0	0	0	4	0	0.4	1	5	7	9	5	4	4	11	5.7	0
558	RED-TAILED BRISTLEBILL <i>Bleda syndactyla</i>	RB	FF		0	3	0	6	2	6	1	0	0	2.2	0	2	6	1	3	3	2	4	2.6	0
559	GREEN-TAILED BRISTLEBILL <i>Bleda eximia</i>	RB	FF		0	0	0	0	0	0	2	0	0	0.8	0	0	4	0	2	1	4	3	1.8	0
561	RED-TAILED GREENBUL <i>Criniger calurus</i>	RB	FF		0	0	0	0	5	5	5	6	6	3.7	0	1	2	0	0	0	0	0	0.4	0
562	COMMON BULBUL <i>Pycnonotus barbatus</i>	RB	f		4	4	4	0	0	0	0	0	0	1.2	0	0	0	0	0	0	0	0	0.0	0
563	WESTERN NICATOR <i>Nicator chloris</i>	RB	F		0	0	0	3	6	0	0	0	5	1.8	0	0	0	0	0	0	0	0	0.0	0
566	FOREST ROBIN <i>Stiphrornis erythrorhox</i>	RB	FF		0	0	0	0	0	0	0	4	0	0.4	0	6	6	4	3	2	2	5	3.5	0
575	BLUE-SHOULDERED ROBIN-CHAT <i>Cossypha cyanocampter</i>	RB	F		0	5	0	0	0	1	2	0	5	1.3	1	0	1	0	1	0	2	0	0.6	0
577	RED-CAPPED ROBIN-CHAT <i>Cossypha natalensis</i>	RB	F		0	0	0	0	0	0	0	0	0	0.0	1	2	0	0	0	0	0	0	0.4	0
579	FIRE-CRESTED ALETHE <i>Alethe diademata</i>	RB	FF		0	0	0	0	0	0	6	0	0	0.6	0	3	3	4	4	4	3	1	2.7	0
581	BROWN-CHESTED ALETHE <i>Alethe poliocephala</i>	RB	FF		0	0	0	0	0	0	0	0	0	0.0	0	1	13	1	2	7	0	2	2.3	0
584	RUFOUS FLYCATCHER-THRUSH <i>Stizorhina fraseri</i>	RB	FF	G-VU	0	4	0	6	0	5	6	0	5	3.7	0	0	0	1	1	0	0	1	0.4	0
670	BLACK-THROATED APALIS <i>Apalis jacksoni</i>	RB	FF		0	0	0	0	0	0	5	5	5	2.2	0	0	0	0	0	0	0	0	0.0	0
677	GREY-BACKED CAMAROPTERA <i>Camaroptera brachyura</i>	RB	f		4	4	0	0	0	2	3	4	1	2.7	1	0	0	0	0	0	0	1	0.3	0
679	OLIVE-GREEN CAMAROPTERA <i>Camaroptera chloronota</i>	RB	FF		0	0	0	0	6	0	0	0	0	0.6	0	0	0	1	2	0	1	1	0.6	0
709	GREEN HYLIA <i>Hylia prasina</i>	RB	F		0	0	0	0	1	0	0	5	0	1.0	1	0	0	0	1	0	1	0	0.4	0
719	ASHY FLYCATCHER <i>Muscicapa caerulescens</i>	RB	F		0	0	0	0	0	0	0	0	0	0.4	0	0	0	0	0	0	0	0	0.0	0
734	DUSKY CRESTED-FLYCATCHER <i>Trochocercus nigromitrata</i>	RB	F		0	0	0	0	0	0	0	0	0	0.0	0	0	1	0	0	0	0	0	0.1	0

Atlas No ^a	Species	Status ^b	Spec ^c	RD ^d	T1	T1	T2	T3	T4	T5	T6	T7	T8	Mean TSC	M1	M2	M3	M4	M5	M6	M7	M8	Mean Net	OP
739	AFRICAN PARADISE-FLYCATCHER <i>Terpsiphone viridis</i>	RB	f		0	0	0	0	0	2	0	0	0	0.2	0	0	0	0	0	0	0	0	0.0	0
740	RED-BELLIED PARADISE-FLYCATCHER <i>Terpsiphone rufiventer</i>	RB	F		0	2	1	0	0	0	6	4	2	1.5	0	1	0	0	0	1	1	0	0.4	0
743	CHESTNUT WATTLE-EYE <i>Dyphorophya castanea</i>	RB	FF		0	2	0	3	6	5	6	2	3	2.7	1	0	0	0	0	0	0	0	0.1	0
744	JAMESON'S WATTLE-EYE <i>Dyphorophya jamesoni</i>	RB	FF		0	0	0	0	0	0	0	0	0	0.0	0	0	0	0	0	0	1	1	0.3	0
755	BROWN ILLADOPSIS <i>Illadopsis fulvescens</i>	RB	FF		0	0	0	1	0	5	0	0	0	1.2	0	1	4	4	3	0	0	3	1.9	0
757	SCALY-BREASTED ILLADOPSIS <i>Illadopsis albipectus</i>	RB	FF		0	3	4	0	0	6	2	0	3	2.4	0	1	1	1	0	1	2	1	0.9	0
784	OLIVE SUNBIRD <i>Cyanomitra olivacea</i>	RB	FF		0	5	5	0	6	6	0	4	0	2.6	1	2	3	4	1	3	6	2	2.7	0
794	COLLARED SUNBIRD <i>Hedydipna collaris</i>	RB	F		0	0	0	0	0	0	0	4	0	0.4	0	0	0	0	0	0	0	0	0.0	0
796	OLIVE-BELLIED SUNBIRD <i>Cinnyris chloropygia</i>	RB	F		0	0	5	0	0	0	0	0	4	1.8	0	0	0	0	0	0	0	0	0.0	0
811	YELLOW WHITE-EYE <i>Zosterops senegalensis</i>	RB	f		0	1	0	0	0	0	0	0	0	1.0	0	0	0	0	0	0	0	0	0.0	0
831	BROWN-CROWNED TCHAGRA <i>Tchagra australis</i>	RB			0	3	0	0	0	0	0	0	0	0.3	0	0	0	0	0	0	0	0	0.0	0
848	WESTERN BLACK-HEADED ORIOLE <i>Oriolus brachyrhynchus</i>	RB	F		0	0	0	5	2	6	6	6	6	4.3	0	0	0	0	0	0	0	0	0.0	0
853	FORK-TAILED DRONGO <i>Dicrurus adsimilis</i>	RB	f/F		0	0	0	0	0	2	0	0	0	0.2	0	0	0	0	0	0	0	0	0.0	0
937	GROSBEAK WEAVER <i>Amblyospiza albifrons</i>	RB	fW		0	0	0	2	0	0	0	0	0	0.2	0	0	0	0	0	0	0	0	0.0	0
954	RED-HEADED BLUEBILL <i>Spermophaga ruficapilla</i>	RB	F		0	0	0	0	0	0	0	0	0	0.0	0	2	0	2	2	0	0	0	0.6	0
976	ZEBRA WAXBILL <i>Amandava subflava</i>	RB			0	0	0	0	0	0	0	0	0	0.0	1	0	2	0	0	0	0	0	0.4	0
	Count Totals ^e			Σ	9	18	16	22	17	26	30	25	27		.									
				new	9	20	4	11	3	4	8	5	1											
				ΣΣ	9	20	24	35	38	42	50	55	55											

Notes:

- a Species numbers and names correspond to the Uganda bird check-list (NBDB, unpubl) and Carswell *et al* (2005) respectively.
- b RB = resident breeding species; R (B) refers to a species whose breeding is probable in Uganda but has not been confirmed; AfM is an Afrotropical migrant, PM a Palearctic migrant and NB means non-breeding. A query mark (?) indicates uncertainty.
- c Specialisations of species are listed in Table B4
- d Red Data species (IUCN, 2006; Bennun and Njoroge, 1996)
- e Σ is the total number of species; these are accumulated by adding those species which were new in successive counts to give the running total (ΣΣ).

Appendix B2. Summary of bird count data from the smaller forests, and from the wayleave in Mabira, which is a non-forest site. There were two TSCs in the Mabira wayleave area (T9, 10); three in Kifu (K1, 2, 3 and the mean KM) and one at Namyoya (N)

Atlas No ^a	Species	Status ^b	Spec ^c	RD ^d	T9	T10	K1	K2	K3	KM	N
26	BLACK-HEADED HERON <i>Ardea melanocephala</i>	RB	w		0	0	0	0	0	0	6
75	BLACK KITE <i>Milvus migrans</i>	RB, PM	pA		0	0	0	0	0	0	6
109	LIZARD BUZZARD <i>Kaupifalco monogrammicus</i>	RB	F		0	3	0	0	0	0	0
142	HELMETED GUINEAFOWL <i>Numida meleagris</i>	RB	G		0	0	0	0	0	0	1
168	WHITE-SPOTTED FLUEFTAIL <i>Sarothrura pulchra</i>	RB	FW		1	4	4	6	0	3.3	0
270	TAMBOURINE DOVE <i>Turtur tympanistria</i>	RB	F		6	6	0	4	0	1.3	0
271	BLUE-SPOTTED WOOD-DOVE <i>Turtur afer</i>	RB	F		4	0	6	6	6	6.0	5
283	RED-EYED DOVE <i>Streptopelia semitorquata</i>	RB	f		0	0	3	0	0	1.0	3
290	GREY PARROT <i>Psittacus erithacus</i>	RB	FF	R-NT	5	0	0	0	0	0	5
296	GREAT BLUE TURACO <i>Corythaeola cristata</i>	RB	F		0	1	4	4	0	2.7	1
297	BLACK-BILLED TURACO <i>Tauraco schuetti</i>	RB	FF		0	5	0	0	0	0	0
302	ROSS'S TURACO <i>Musophaga rossae</i>	RB	F		0	0	6	0	0	2.0	0
305	EASTERN GREY PLANTAIN-EATER <i>Crinifer zonurus</i>	RB			0	0	6	0	0	2.0	2
309	RED-CHESTED CUCKOO <i>Cuculus solitarius</i>	RB, AfM/NB?	AF		0	0	5	5	0	3.3	5
314	DUSKY LONG-TAILED CUCKOO <i>Cercococcyx mechowi</i>	RB	FF		0	4	0	0	0	0	0
317	AFRICAN EMERALD CUCKOO <i>Chrysococcyx cupreus</i>	RB?	F		2	6	4	2	4	3.3	0
319	KLAAS' CUCKOO <i>Chrysococcyx klaas</i>	RB	f		0	5	4	0	0	1.3	4
320	DIEDERIK CUCKOO <i>Chrysococcyx caprius</i>	RB, AfM/(B)? PM			0	0	0	0	3	1.0	0
321	YELLOWBILL <i>Ceuthmochares aereus</i>	RB	F		0	1	0	0	0	0	0
323	WHITE-BROWED COUCAL <i>Centropus superciliosus</i>	RB			6	0	5	0	4	3.0	1
358	AFRICAN PALM SWIFT <i>Cypsiurus parvus</i>	RB			1	0	0	0	0	0	0
367	ALPINE SWIFT <i>Apus melba</i>	RB	p		4	0	0	0	0	0	0
369	SPECKLED MOUSEBIRD <i>Colius striatus</i>	RB			0	0	0	1	4	1.7	4
371	NARINA'S TROGON <i>Apaloderma narina</i>	RB	F		1	6	0	0	0	0	0
375	WOODLAND KINGFISHER <i>Halcyon senegalensis</i>	PM, RB	A		0	0	5	0	3	2.7	0
378	AFRICAN PYGMY KINGFISHER <i>Ceyx picta</i>	RB, AfM/NB	fw		0	0	0	0	0	0	4
390	WHITE-THROATED BEE-EATER <i>Merops albicollis</i>	AfM/NB, FB, PM	Af		0	1	0	0	0	0	0
401	BROAD-BILLED ROLLER <i>Eurystomus glaucurus</i>	RB, AfM/NB?	Afw		0	0	6	1	6	4.3	0
419	CROWNED HORNBILL <i>Tockus alboterminatus</i>	RB	f		0	0	5	0	0	1.7	0
422	BLACK-AND-WHITE CASQUED HORNBILL <i>Bycanistes subcylindricus</i>	RB	F		5	6	6	3	6	5.0	6
426	SPECKLED TINKERBIRD <i>Pogoniulus scolopaceus</i>	RB	F		0	6	0	2	5	2.3	0
430	YELLOW-THROATED TINKERBIRD <i>Pogoniulus subsulphureus</i>	RB	FF		5	6	6	1	6	4.3	0
431	YELLOW-RUMPED TINKERBIRD <i>Pogoniulus bilineatus</i>	RB	F		6	6	5	0	4	3.0	6

Atlas No ^a	Species	Status ^b	Spec ^c	RD ^d	T9	T10	K1	K2	K3	KM	N
434	YELLOW-SPOTTED BARBET <i>Buccanodon duchaillui</i>	RB	FF		6	0	0	0	6	2.0	0
435	HAIRY-BREASTED BARBET <i>Tricholaema hirsuta</i>	RB	F		1	0	0	0	0	0	0
445	YELLOW-BILLED BARBET <i>Trachyphonus purpuratus</i>	RB	FF		1	6	3	0	0	1.0	0
456	LESSER HONEYGUIDE <i>Indicator minor</i>	RB	f		0	0	0	5	0	1.7	0
498	WHITE-HEADED SAW-WING <i>Psalidoprocne albiceps</i>	RB, AfM/NB?	f	R-RR	0	3	0	2	6	2.7	0
505	LESSER STRIPED SWALLOW <i>Hirundo abyssinica</i>	RB			0	0	0	0	1	0.3	0
538	LITTLE GREENBUL <i>Andropadus virens</i>	RB	F		5	6	6	6	6	6.0	0
542	YELLOW-WHISKERED GREENBUL <i>Andropadus latirostris</i>	RB	F		6	0	0	0	0	0	0
558	RED-TAILED BRISTLEBILL <i>Bleda syndactyla</i>	RB	FF		4	0	0	0	0	0	0
559	GREEN-TAILED BRISTLEBILL <i>Bleda eximia</i>	RB	FF		0	6	0	0	0	0	0
561	RED-TAILED GREENBUL <i>Criniger calurus</i>	RB	FF		0	5	0	0	0	0	0
562	COMMON BULBUL <i>Pycnonotus barbatus</i>	RB	f		0	0	0	5	4	3.0	5
563	WESTERN NICATOR <i>Nicator chloris</i>	RB	F		2	2	0	0	0	0	0
584	RUFIOUS FLYCATCHER-THRUSH <i>Stizorhina fraseri</i>	RB	FF	G-VU	5	6	0	0	0	0	0
638	RED-FACED CISTICOLA <i>Cisticola erythrops</i>	RB	w		0	0	0	0	0	0	3
647	WINDING CISTICOLA <i>Cisticola galactotes</i>	RB	w		0	0	0	0	0	0	2
658	TAWNY-FLANKED PRINIA <i>Prinia subflava</i>	RB	fw		0	0	0	0	0	0	5
662	WHITE-CHINNED PRINIA <i>Prinia leucopogon</i>	RB	F		0	0	1	5	0	2.0	0
670	BLACK-THROATED APALIS <i>Apalis jacksoni</i>	RB	FF		2	5	0	0	0	0	0
677	GREY-BACKED CAMAROPTERA <i>Camaroptera brachyura</i>	RB	f		4	5	0	5	4	3.0	5
709	GREEN HYLIA <i>Hylia prasina</i>	RB	F		4	0	0	3	2	1.7	0
719	ASHY FLYCATCHER <i>Muscicapa caerulescens</i>	RB	F		4	0	0	0	0	0	0
742	BLACK-AND-WHITE SHRIKE-FLYCATCHER <i>Bias musicus</i>	RB	f		0	0	0	0	1	0.3	0
746	BROWN-THROATED WATTLE-EYE <i>Platysteira cyanea</i>	RB	f		0	0	0	6	0	2.0	2
755	BROWN ILLADOPSIS <i>Illadopsis fulvescens</i>	RB	FF		6	0	0	0	0	0	0
757	SCALY-BREASTED ILLADOPSIS <i>Illadopsis albipectus</i>	RB	FF		6	0	0	0	0	0	0
796	OLIVE-BELLIED SUNBIRD <i>Cinnyris chloropygia</i>	RB	F		4	5	0	0	4	1.3	5
809	SUPERB SUNBIRD <i>Cinnyris superba</i>	RB	F		0	0	0	4	0	1.3	0
810	COPPER SUNBIRD <i>Cinnyris cuprea</i>	RB	fw		0	0	0	0	0	0	3
831	BROWN-CROWNED TCHAGRA <i>Tchagra australis</i>	RB			0	0	0	0	0	0	3
848	WESTERN BLACK-HEADED ORIOLE <i>Oriolus brachyrhynchus</i>	RB	F		6	6	0	0	0	0	0
871	SPLENDID GLOSSY STARLING <i>Lamprotornis splendidus</i>	AfM/NB? RB	F		0	0	2	1	4	2.3	6
872	RÜPPELL'S LONG-TAILED STARLING <i>Lamprotornis purpuropterus</i>	RB			0	0	2	0	0	0.7	0
893	BAGLAFECHT WEAVER <i>Ploceus baglafecht</i>	RB	f		0	0	0	0	0	0	2
896	BLACK-NECKED WEAVER <i>Ploceus nigricollis</i>	RB	f		0	0	0	0	0	0	1

Atlas No ^a	Species	Status ^b	Spec ^c	RD ^d	T9	T10	K1	K2	K3	KM	N
897	SPECTACLED WEAVER <i>Ploceus ocularis</i>	RB	f		0	0	0	0	1	0.3	0
907	VIEILLOT'S BLACK WEAVER <i>Ploceus nigerrimus</i>	RB	f		0	0	0	0	0	0	4
908	BLACK-HEADED WEAVER <i>Ploceus cucullatus</i>	RB			0	0	0	1	0	0.3	0
913	YELLOW-MANTLED WEAVER <i>Ploceus tricolor</i>	RB	FF		0	0	0	1	0	0.3	0
932	FAN-TAILED WIDOWBIRD <i>Euplectes axillaris</i>	RB	w		0	0	0	0	0	0	1
937	GROSBEAK WEAVER <i>Amblyospiza albifrons</i>	RB	fW		0	0	0	0	0	0	3
942	WHITE-BREASTED NEGROFINCH <i>Nigrita fusconota</i>	RB	F		0	0	0	0	1	0.3	0
970	BLACK-CROWNED WAXBILL <i>Estrilda nonnula</i>	RB	f		0	0	0	0	0	0	3
980	BRONZE MANNIKIN <i>Lonchura cucullata</i>	RB			0	0	0	0	4	1.3	5
981	BLACK-AND-WHITE MANNIKIN <i>Lonchura bicolor</i>	RB	f		0	0	0	0	0	0	3
995	YELLOW-FRONTED CANARY <i>Serinus mozambicus</i>	RB			0	0	0	0	0	0	3

Notes a to d: see Appendix B1

Appendix B3. Bird records from Lubigi wetland

Atlas ^a No.	Species	Status ^b	Sp ^c	RD ^d	TSC scores						
					Site1	Site2	Site 3	Site 4	Site 5	Site 6	Mean
9	PINK-BACKED PELICAN <i>Pelecanus rufescens</i>	RB	W		5						0.8
14	COMMON SQUACCO HERON <i>Ardeola ralloides</i>	WV?, AfM/NB?, RB	W						4	3	1.2
17	CATTLE EGRET <i>Bubulcus ibis</i>	RB	G		5	5			4	2	2.7
21	LITTLE EGRET <i>Egretta garzetta</i>	RB	W							4	0.7
23	GREAT [=WHITE] EGRET <i>Casmerodius alba</i>	RN?, AfM/NB?	W	R-VU						5	3.8
24	PURPLE HERON <i>Ardea purpurea</i>	RB? FB	W	R-NT						1	0.2
25	GREY HERON <i>Ardea cinerea</i>	RB? FB, OW?	W	R-NT						4	0.7
26	BLACK-HEADED HERON <i>Ardea melanocephala</i>	RB	w							4	0.7
28	HAMERKOP <i>Scopus umbretta</i>	RB	w		6	6	3			6	3.5
36	MARABOU STORK <i>Leptoptilos crumeniferus</i>	RB, AfM/B	w		4	6	6	3			3.2
39	HADADA <i>Bostrychia hagedash</i>	RB	w		5					1	1.0
75	BLACK KITE <i>Milvus migrans</i>	RB, PM	pA							2	0.3
80	HOODED VULTURE <i>Necrosyrtes monachus</i>	RB	f		5	1	6	3			2.5
90	AFRICAN HARRIER-HAWK <i>Polyboroides typus</i>	RB	f						1		0.2
93	AFRICAN MARSH HARRIER <i>Circus ranivorus</i>	R(B)	W	R-NT						5	0.8
109	LIZARD BUZZARD <i>Kaupifalco monogrammicus</i>	RB	F						6		1.0
117	WAHLBERG'S EAGLE <i>Aquila wahlbergi</i>	AfM/NB, RB	Af						1		0.2
132	GREY KESTREL <i>Falco ardosiaceus</i>	RB							1		0.2
178	BLACK CRAKE <i>Amaurornis flavirostris</i>	RB	W			5				1	1.0
185	GREY CROWNED CRANE <i>Balearica regulorum</i>	RB, AfM/NB?	WG	R-NT						1	0.2
221	AFRICAN WATTLED LAPWING [=PLOVER] <i>Vanellus senegallus</i>	RB	W		3						0.5
268	AFRICAN GREEN-PIGEON <i>Treron calva</i>	RB	F				2				0.3
270	TAMBOURINE DOVE <i>Turtur tympanistria</i>	RB	F					2	2	1	0.8
271	BLUE-SPOTTED WOOD-DOVE <i>Turtur afer</i>	RB	F		4			2		1	1.2
283	RED-EYED DOVE <i>Streptopelia semitorquata</i>	RB	f		6		5		5	2	3.0
289	LAUGHING DOVE <i>Streptopelia senegalensis</i>	RB			5		1		2		1.3
293	RED-HEADED LOVEBIRD <i>Agapornis pullaria</i>	AfM/NB? RB	F				1	6			1.2
305	EASTERN GREY PLANTAIN-EATER <i>Crinifer zonurus</i>	RB							6	6	2.0
317	AFRICAN EMERALD CUCKOO <i>Chrysococcyx cupreus</i>	RB?	F						5		0.8
319	KLAAS' CUCKOO <i>Chrysococcyx klaas</i>	RB	f			1					0.2
320	DIEDERIK [=DIDRIC] CUCKOO <i>Chrysococcyx caprius</i>	RB, AfM/(B)? PM							3		0.5

Atlas ^a No.	Species	Status ^b	Spc ^c	RD ^d	TSC scores						
					Site1	Site2	Site 3	Site 4	Site 5	Site 6	Mean
326	BLUE-HEADED COUCAL <i>Centropus monachus</i>	RB	W		1		2			1	0.7
358	AFRICAN PALM SWIFT <i>Cypsiurus parvus</i>	RB						1			0.2
369	SPECKLED MOUSEBIRD <i>Colius striatus</i>	RB			6			4	6		2.7
375	WOODLAND KINGFISHER <i>Halcyon senegalensis</i>	PM, RB								1	0.2
385	LITTLE BEE-EATER <i>Merops pusillus</i>	RB	G						6		1.0
386	BLUE-BREASTED BEE-EATER <i>Merops variegatus</i>	RB	W						2		0.3
390	WHITE-THROATED BEE-EATER <i>Merops albicollis</i>	AfM/NB, FB, PM	Af		6		6	5			2.8
392	BLUE-CHEEKED BEE-EATER <i>Merops persicus</i>	WV, PM	P					5			0.8
401	BROAD-BILLED ROLLER <i>Eurystomus glaucurus</i>	RB, AfM/NB?	Afw		5	4	3	6	6		4.0
419	CROWNED HORNBILL <i>Tockus alboterminatus</i>	RB	f						5		0.8
433	YELLOW-FRONTED TINKERBIRD <i>Pogoniulus chrysoconus</i>	RB	f				1				0.2
443	DOUBLE-TOOTHED BARBET <i>Lybius bidentatus</i>	RB	f		5				6		1.8
465	NUBIAN WOODPECKER <i>Campethera nubica</i>	RB					5			1	1.0
498	WHITE-HEADED SAW-WING [=ROUGHWING] <i>Psalidoprocne albiceps</i>	RB, AfM/NB?	f	R-RR				1		2	0.5
512	ANGOLA SWALLOW <i>Hirundo angolensis</i>	RB, AfM/B?	w			3					0.5
520	AFRICAN PIED WAGTAIL <i>Motacilla aguimp</i>	RB	w							2	0.3
529	YELLOW-THROATED LONGCLAW <i>Macronyx croceus</i>	RB	G		6		1	6			2.2
538	LITTLE GREENBUL <i>Andropadus virens</i>	RB	F				6	4	2		2.0
547	YELLOW-THROATED GREENBUL [=LEAFLOVE] <i>Chlorocichla flavicollis</i>	RB	f				6				1.0
562	COMMON BULBUL <i>Pycnonotus barbatus</i>	RB	f		5	6	6		6	2	4.2
576	WHITE-BROWED ROBIN-CHAT <i>Cossypha heuglini</i>	RB	f				6				1.0
615	WHITE-WINGED WARBLER <i>Bradypterus carpalis</i>	R(B)	E, W		5		9		1	5	2.8
621	AFRICAN MOUSTACHED WARBLER <i>Melocichla mentalis</i>	RB							4		0.7
630	GREATER SWAMP WARBLER <i>Acrocephalus rufescens</i>	RB	e, W		6	6	2		4	5	0.8
631	LESSER SWAMP WARBLER <i>Acrocephalus gracilirostris</i>	RB	w							1	0.2
638	RED-FACED CISTICOLA <i>Cisticola erythrops</i>	RB	w		5				6		1.8
647	WINDING CISTICOLA <i>Cisticola galactotes</i>	RB	w		6	6	6		5	6	4.8
648	CARRUTHERS'S CISTICOLA <i>Cisticola carruthersi</i>	RB	E, W	R-RR	5		4	6		6	3.5
658	TAWNY-FLANKED PRINIA <i>Prinia subflava</i>	RB	fw				5				0.8
662	WHITE-CHINNED PRINIA <i>Prinia leucopogon</i>	RB	F		5		6				1.8
677	GREY-BACKED CAMAROPTERA <i>Camaroptera brachyura</i>	RB	f						6	1	1.2
701	GREY-CAPPED WARBLER <i>Eminia lepida</i>	RB	fw	R-RR	3	5	6	6	1	6	4.5
720	SWAMP FLYCATCHER <i>Muscicapa aquatica</i>	RB	W		1					2	0.5

Atlas ^a No.	Species	Status ^b	Spc ^c	RD ^d	TSC scores						
					Site1	Site2	Site 3	Site 4	Site 5	Site 6	Mean
746	BROWN-THROATED WATTLE-EYE <i>Platysteira cyanea</i>	RB	f				5				0.8
764	BLACK-LORED BABBLER <i>Turdoides sharpei</i>	RB		R-RR	6						1.0
781	GREEN-HEADED SUNBIRD <i>Cyanomitra verticalis</i>	RB	F		3					4	1.2
784	OLIVE SUNBIRD <i>Cyanomitra olivacea</i>	RB	FF		2	3		6			1.8
787	SCARLET-CHESTED SUNBIRD <i>Chalcomitra senegalensis</i>	RB	f				6	5		5	2.7
802	MARICO [=MARIQUA] SUNBIRD <i>Cinnyris mariquensis</i>	RB			2			5			1.2
803	RED-CHESTED SUNBIRD <i>Cinnyris erythrocerca</i>	RB	W	R-RR	5				6	6	2.8
810	COPPER SUNBIRD <i>Cinnyris cuprea</i>	RB	fw			6		6			2.0
815	GREY-BACKED FISCAL <i>Lanius excubitoroides</i>	RB	Afw			6					1.0
828	SULPHUR-BREASTED BUSH-SHRIKE <i>Malaconotus sulfureopectus</i>	RB? AfM/B?	f						4		0.7
830	MARSH TCHAGRA <i>Tchagra minutus</i>	RB	w						2		0.3
842	PAPYRUS GONOLEK <i>Laniarius mufumbiri</i>	R(B)	E, w		6	6	2		6	6	4.3
843	BLACK-HEADED GONOLEK <i>Laniarius erythrogaster</i>	RB	f				6		5		1.8
855	PIED CROW <i>Corvus albus</i>	RB			4	2				2	1.3
872	RÜPPELL'S LONG-TAILED [=GLOSSY] STARLING <i>Lamprolornis purpuropterus</i>	RB			1	4	1	6			2.0
881	GREY-HEADED SPARROW <i>Passer griseus</i>	RB				3				2	0.8
894	SLENDER-BILLED WEAVER <i>Ploceus pelzelni</i>	RB	fW		4	5			5	3	2.8
897	SPECTACLED WEAVER <i>Ploceus ocularis</i>	RB	f				6				1.0
908	BLACK-HEADED WEAVER <i>Ploceus cucullatus</i>	RB								3	0.5
910	YELLOW-BACKED WEAVER <i>Ploceus melanocephalus</i>	RB	W						4	1	0.8
911	GOLDEN-BACKED WEAVER <i>Ploceus jacksoni</i>	RB	w	R-RR			6				1.0
915	COMPACT WEAVER <i>Ploceus superciliosus</i>	RB	fw				5		1		1.0
932	FAN-TAILED WIDOWBIRD <i>Euplectes axillaris</i>	RB	w				6	6	6	2	3.3
943	WHITE-COLLARED OLIVEBACK <i>Nesocharis ansorgei</i>	R(B)	fw	R-RR					1		0.2
959	RED-BILLED FIREFINCH <i>Lagonosticta senegala</i>	RB					2				0.3
969	COMMON WAXBILL <i>Estrilda astrild</i>	RB	wG		5	6	5	4			3.3
980	BRONZE MANNIKIN <i>Lonchura cucullata</i>	RB			2	6	6		1	6	3.5
981	BLACK-AND-WHITE MANNIKIN <i>Lonchura bicolor</i>	RB	f		1		5	4			1.7
995	YELLOW-FRONTED CANARY <i>Serinus mozambicus</i>	RB				1				1	0.3
			Σ		37	23	37	23	38	43	

Notes: a to d are as for Appendix B1.

Appendix M1. Mammal species recorded along the various transects and locations surveyed in Mabira Forest.

Species	Site 1	Site 2	Site 3	Site 4	Site 5	Site 6	Site 7	Site 8	General Wayleave
Insectivora									
Uganda forest Musk Shrew (<i>Crocidura selina</i>)		√		√					
Northern Giant Musk Shrew (<i>Crocidura olivieri</i>)								√	
Hero Shrew (<i>Scutisorex somereni</i>)					√	√	√	√	
Chiroptera									
Straw colored Fruit Bat (<i>Eidolon helvum</i>)	√	√						√	
Little epauletted Fruit Bat (<i>Epomophorus labiatus</i>)		√						√	√
African Long-tongued Fruit Bat (<i>Megaloglossus woermanni</i>)									√
Bocage's Fruit Bat (<i>Rousettus angolensis</i>)			√		√		√	√	
Noack's Leaf-nosed Bat (<i>Hipposideros ruber</i>)					√				
Halcyon Horseshoe Bat (<i>Rhinolophus alcyone</i>)		√	√		√			√	
Banana Bat (<i>Pipistrellus nanus</i>)									√
Primates									
Red tailed Monkey (<i>Cercopithecus ascanius</i>)	√	√	√	√	√	√	√	√	
Galago (<i>Galago senegalensis</i>)		√							
Grey Cheeked Mangabey (<i>Cercocebus abigena</i>)							√		
Carnivora									
Marsh Mongoose (<i>Atilax paludinosus</i>)									√
Forest Genet (<i>Genetta victoriae</i>)					√				
Dwarf Mongoose (<i>Hologale parvula</i>)						√			√
Slender Mongoose (<i>Herpestes ichneumon</i>)							√		
Side Striped Jackal (<i>Canis adustus</i>)									√
Serval Cat (<i>Felis serval</i>)									√
Pholidota									
Tree Pangolin (<i>Manis tricuspis</i>)				√					
Hyracoidea									
Tree Hyrax (<i>Dendrohyrax aboreaus</i>)	√	√	√	√	√	√	√	√	
Artiodactyla									
Blue Duiker (<i>Cephalophus monticola</i>)			√	√					
Bushpig (<i>Potamochoerus porcus</i>)			√						
Red Forest Duiker (<i>Cephalophus harveyi</i>)			√						
Bushbuck (<i>Tragelaphus scriptus</i>)			√						
Rodentia									
Congo forest Rat (<i>Deomys ferugineous</i>)					√				
Stella Wood Mouse (<i>Hylomyscus stella</i>)	√		√	√		√	√		
Eastern B rush-furred Mouse (<i>Lophuromys flavopunctatus</i>)			√					√	
Common Brush furred Mouse (<i>Lophuromys sikapusi</i>)			√						
Peter's Stripped Mouse (<i>Hybomys univittatus</i>)			√		√				
Long footed rat (<i>Malacomys longipes</i>)			√						
Jackson's Soft-furred Rat (<i>Praomys jacksoni</i>)			√		√	√	√	√	
Stripped Ground Squirrel (<i>Xerus erythropus</i>)					√				
Brush tailed Porcupine (<i>Atherurus africanus</i>)			√						
Macroscelidea									
Giant Elephant Shrew (<i>Rhynchocyon cirnei</i>)		√							
Totals	4	8	15	6	9	8	8	10	7



Plate 1: Some of the bigger diameter trees that are found in the sites 5, 6,7 and 8 (Cluster M1 of Figure P3) of Mabira forest. *Dracaena fragrans*, a common forest floor shrub is in the foreground.



Plate 2: A *Broussonetia papyrifera* dominated stand. This species is characteristic of study Sites 1 and 2 in Mabira forest.



a)
Garden of Cassava either side of the transmission line in Mabira (a & b)



b)



c) Newly tilled garden



d) Maize garden in one section along the line

Plate 3(a- d): Various scenes of cultivation along the transmission line in Mabira forest towards Wasswa Village



a)



b)

Plate 4: Transmission line rising over the canopy (a & b) at the low points in the forest



Plate 5: Dense growth of vegetation along wayleave in Kifu



Plate 6: Regeneration in Kifu Forest within *Maesopsis eminii*



Plate 7: A plantation of *Auracaria cunninghamii* on the immediate northern side of the existing power line in Kifu forest reserve.



Plate 8: A variety of ongoing/existent human impacts that we recorded in Lubigi Wetlands (a – h)



a. Cultivation



b. Hollows left after mining sand or clay



c. An area Quarried for rocks



d. Biodegradable Refuse dumping



e. Non bio-degradable refuse



f. Non bio degradable after its burnt



g. A variety of domestic refuse



h. Papyrus harvesting



Plate 9: A transect located through a thick *papyrus* stand at Site 1



Plate 10: Another human impact Block *Cyperus* Making in Lubigi

Plate 11: Different vegetation communities that characterise the Lubigi Swamp



a. *Phoenix reclinata* and *Alchornea cordifolia* dominated swamp fringes



b. *Cyperus* and *Scleria* dominated seasonally flooded grassland



c. *Aeschynomene indica* dominated swamp fringe



d. *Papyrus* & *Phoenix* co-dominated Swamp fringes

Plate 12: Situational photos taken in Namyoya/ Mwola Forest



a) Pylon 51 facing towards the tea plantations



b) *Eucalyptus* plantations in the fore ground



c) Maize growing in the *Eucalyptus*



d) Field of harvested *Eucalyptus*



e) Field of harvested *Eucalyptus*



f) Field of harvested *Eucalyptus*




Plate 13: The *Eucalyptus* plantation on the immediate northern side of the existing power line in Namyoya forest reserve: (a) the plantation landscape view (b) the *Eucalyptus* stand



Appendix B.4

**Excerpts from NFA's Forest Nature
Conservation Plan, March 1999**



APPENDIX 16: MABIRA FOREST PROFILE

(Category: CORE conservation forest)

1 Basis for selection

The forest was selected for Nature Reserve establishment in recognition of its biodiversity importance, especially because:

- the site contributes more than 1% of the national protected area system complement.
- the site supports at least one unique species of tree of conservation importance.
- the site supports vegetation type D1(Langdale - Brown et al., 1964) otherwise not represented in protected area system of Uganda.

2 Physical description

Area and demarcation: Area: 306 km²; (311 km², with Namananga/Namawanyi); with a total boundary length 347.4 km, all adjoining community lands. The boundary is largely artificial, maintained as a cut-line with corner cairns and directional trenches.

Establishment: First established, under the Buganda Agreement (1900) and later formally gazetted in 1932.

Location: On the Kampala-Jinja highway at about 54 km from Kampala and 26 km from Jinja, 20 km north of the Lake Victoria shoreline, in central Uganda, between 0°24'-0°35' N and 32°52'-33°07' E. The reserve occupies part of counties of Ntenjeru, Nakifuma, Buikwe, and Mukono, all in Mukono District. Covered by Uganda Department of Lands and Surveys map sheets 61/4, 62/3, 71/2 and 72/1 at 1:50,000.

Physical features: The reserve occupies gently undulating terrain with numerous flat-topped hills, with altitudes of 1070-1340 m a.s.l., with less than 10% exceeding 5° slope. The area is drained by two main rivers, the Musamya and Sezibwa, which flow northwards into Lake Kyoga.

3 Vegetation and forest condition

The majority of the area (292km²; 95%) is occupied by Tropical High Forest communities, classified as type D1 (*Celtis-Chrysophyllum* medium altitude moist semi-deciduous forest) and the remainder (5%) by *Piptadeniastrum-Albizia-Celtis* medium altitude moist evergreen forest (Langdale - Brown et al., 1964). Human activities have greatly influenced the forest condition, making some areas characteristic sub-climaxes. Sub-types of vegetation present are young or colonising mixed forest, dominated by *Maesopsis eminii* (25%), young mixed *Celtis holoptelea* (60%), and mixed forest of wet valley bottoms dominated by *Baikaea insignis*.

A detailed forest type map is available at Forest Department headquarters, based on the 1950s aerial photography, and also reproduced in Howard (1991).

The forest is largely disturbed by human activity (overall condition score 2), mainly because it is located between the two largest urban centres in Uganda, and the area is largely accessible. There has been extensive pitsawing activity and agricultural encroachment (1973-1987). Hunting is widespread.

Forest integrity scores: Settlement = 2, Hunting = 2 Livestock = 1; Timber = 4 Fire = 0; Community = 3 Mining = 1.

4 Economic importance

Community use values: The forest is situated in one of the most densely populated parts of the country (235 people per km² in 1991). Pressure on the peripheral areas of the forest for firewood, building poles and non-timber forest products is correspondingly high. The forest is largely accessible because of the presence of village enclaves and roads leading to them. The 'community-use' value of the reserve is 20, and it is thus potentially very

important economically to the communities around, and for the two nearest towns; Jinja and Kampala.

Timber production: The forest is an important source of pitsawn timber, providing a registered annual off take of about 4,284m³ of sawn timber over the period 1994-96 (Table 16.1), as well as large volumes of illegally cut timber. A timber inventory by Forest Department (in 1992) provided an estimate of 60m³ per ha. standing volumes of merchantable timber exceeding 50 cm dbh. Records showing the number of registered pitsawyers do not exist. However, timber volumes over the period 1964-1996 are indicated in Table 16.1.

Table 16.1 Timber production in Mabira: 1964-1996

Period	Sawmill	Volume (m ³)
1964-1974	Sick Sawmill & Ginnors Ltd.	15,694
1973-1980	Kiira Sawmill & Plywood Factory	16,321
1981-1989	Kiira Sawmill & Plywood Factory	19,041
1990-1993	Kiira Sawmill & Plywood Factory	-
1994-1996 (July)	Nile Plywood (U) Ltd	2,907
1994-1996 (July)	Jinja Construction and Joinery Ltd.	1,377
Total		55,340

Nadagi compartment (479 ha) has been put aside for the establishment of eucalyptus plantations with temporary permits being issued to potential farmers, and there is potential for expansion of this programme.

Other economic values: The reserve has been locally important as a source of building poles, firewood and medicinal compounds. It has also been important for the production of charcoal. It is located between two of the major urban centres in Uganda, and has potential for ecotourism development based on such attractions as the luxurious flora and fauna, and the scenic rivers Musamya and Sezibwa (on which falls are located). The reserve is important for biodiversity (see below) and thus offers scope for the development of a research and education role.

5 Biodiversity values

Of the 65 forest reserves investigated for biodiversity, Mabira does not score among the highest in terms of overall biodiversity, ranking 24th (score =13.1), but ranks 19th in terms of the rarity value of species represented. The forest supports 9 species found in no other Ugandan forest (including 6 butterflies, 1 moth, 1 bird and 1 tree) and one species endemic to Uganda (Table 16.4). It presents the only block of medium altitude moist semi-deciduous forest type D1 (Langdale-Brown et al., 1964) in the protected area system, a vegetation type that does not occur in any of the country's National Parks or Wildlife Reserves.

6 Present management

The reserve is managed from Lwankima Forest Station, by a Forest Officer. The Mukono District forest office plays a supervisory role. Table 16.2 shows the staffing position for Mabira Forest Reserve. There are three Forest Officers, stationed at Lwankima, Maligita and Najjembe. The one at Najjembe works specifically on tourism development. In addition, a total of 3 Assistant Forest Officers, 8 Forest Rangers and 9 Forest Guards assist in the management of this important forest, and are based at various forest stations as indicated in Table 16.2.

The department has six staff houses at Lwankima Forest Station, the local headquarters of the reserve, and has endeavoured to offer ample housing at all the 12 forest stations on this reserve as indicated in Table 16.3.

Table 16.2 Existing and proposed staff deployment at Mabira forest

Station	Existing (proposed) number of staff by category					Total
	FO	AFO	F.R	F.G	PM	
Lwankima	1 (0)	1 (0)	1* (1)	1* (2)	4 (0)	8 (3)
Maligita	1 (0)	0 (0)	0 (1)	1 (0)	2 (0)	4 (1)
Namawanyi	0 (0)	0 (0)	0 (0)	1 (0)	1 (1)	2 (1)
Naluvule	0 (0)	0 (0)	0 (0)	0 (1)	1 (0)	1 (1)
Kyabana	0 (0)	0 (0)	1 (0)	1* (0)	1 (0)	3 (0)
Buwoola	0 (0)	0 (0)	1 (2)	1 (0)	1 (0)	3 (2)
Najjembe	1 (0)	0 (0)	1* (0)	1 (0)	0 (1)	3 (1)
Wanende	0 (0)	0 (0)	1* (0)	0 (1)	4 (0)	5 (1)
Nandagi	0 (0)	1 (0)	1* (0)	0 (1)	0 (1)	2 (2)
Nagojje	0 (0)	1 (0)	1* (0)	1 (0)	2 (0)	5 (0)
Namulaba	0 (0)	0 (0)	1 (0)	1 (0)	1 (1)	3 (1)
Nazigo	0 (0)	0 (0)	0 (0)	1 (0)	0 (2)	1 (2)
Total	3 (0)	3 (0)	3 + 5* (4)	7 + 2* (5)	17 (6)	40(15)

Note: FO - Forest Officer FG - Forest Guard AFO - Assistant Forest Officer PM - Patrol Man
FR - Forest Ranger, * denotes temporary employment on EU Project, not Government employee.

Table 16.3 shows the status of housing in Mabira Forest Reserve and the proposed requirements in order to offer accommodation to all staff.

Table 16.3 Existing (proposed) staff housing

Station	FD old houses.	FD detached	FD semi	Uniport	Total
Lwankima	6(0)	0(0)	0(1)	2(0)	8(1)
Najjembe	1(0)	1(0)	1(1)	0(0)	3(1)
Wanende	0(0)	1(0)	2(0)	0(0)	3(0)
Buwoola	0(0)	0(0)	1(0)	0(0)	1(0)
Kyabaana	0(0)	1(0)	1(0)	0(0)	2(0)
Maligita	1(0)	1(0)	1(1)	4(0)	7(1)
Naluvule	0(0)	0(0)	1(0)	0(0)	1(0)
Namawanyi	0(0)	0(0)	1(0)	0(0)	1(0)
Nandagi	0(0)	0(0)	1(1)	2(0)	3(1)
Nazigo	1(0)	0(0)	0(0)	0(0)	1(0)
Namulaba	0(0)	1(1)	2(0)	0(0)	3(1)
Nagojje	5(0)	0(0)	0(0)	0(0)	5(0)
	14(0)	5(1)	11(4)	8(0)	38(5)

There are no bicycles or motorcycles to facilitate the management of the forest, inspite of the availability of a road network in the forest reserve. The latest (Interim) Management Plan covered the period (1994-1995) and prescribed for the conservation of the forest biodiversity, the protection of the area's important water catchment role and the maximum yield of hardwood timber. Although a Nature Reserve was proposed, actual demarcation did not take place and discussions were still going on for further changes to the zones. A detailed management plan to cover the period 1997-2007 is now under preparation.

In recent years (since 1990), with the support of the EU-financed Natural Forest Management and Conservation Project, some parts of the boundary have been redemarcated and few sections planted with live markers (see Fig. A16.1). An ecotourism project has also been established and further tourism development is expected.

7 Proposed zonation

Figure A16.1 shows the proposed zonation of the reserve, with one Nature Reserve (approximately 73 km²) one protection zone (approximately 30 km²), recreation zone (approximately 40 km²) and the rest of the reserve (Approximately 170 km²) as production zones.

The proposed Nature Reserve

It will cover the central portion of the forest reserve which is relatively intact. This has been selected to protect a viable area of semi-deciduous forest type D1 (Langdale Brown et. al., 1964), which is important because this is the only protected area in the country in which this forest type is represented.

The proposed protection zone

This will cover the area adjacent to the Nature Reserve with the aim of enhancing the long term viability of the latter. The proposed recreation zone is expected to centre around Najjembe (to the South) and around Musamya river (to the north, near the boundary). The zone encompasses the river and marshes called Musamya, which are a valuable habitat for a number of species of plants and animals, and are some of the most scenic areas of the forest.

The proposed production zones

These cover the majority of the reserve, including the areas that have already been heavily exploited by pitsawyers, the more accessible peripheral areas of the reserve; and most of the south-central parts of the reserve which adjoin a number of enclaves and are more suitable for timber production.

8 Proposed management programme

Staffing: The present staff is inadequate, and redeployment may also be necessary. Most areas lack forest workers; only patrolmen and Forest Guards occur, resulting in inadequate control, and not much labour work on the ground such as planting and boundary maintenance. The Forest Officer at Maligita does not have a Ranger to assist him in his duties. Furthermore, the Forest Guards at the various stations do not have properly motivated and facilitated patrolmen under them. Each guard would need at least four workers and two patrolmen to assist him/her.

Transport will be required as follows: 1-4 wheel drive vehicle for the FO and 3 motorcycles; 1 for Maligita to facilitate operation on the Eastern axis and another for Nagojje for the western part of the reserve, and finally, one for Lwankima forest station. Each Forest Ranger and FG should be facilitated with bicycles. The FO tourism needs to be facilitated with a 4-wheel drive vehicle to enable community outreach programmes, and the running of the visitor's centre. The Forest Officers in charge should be facilitated with transport to carry out effective patrols of the reserve as well as with a radio communication system.

Infrastructure: Four houses will be required for staff, at Lwankima (1 duplex), Najjembe (1 duplex), Namulaba (1 replacement), Maligita (1 duplex) and Nandagi (1 duplex). Details are indicated in Table 16.3.

Demarcation: Over 250 km of reopened external boundary lacks maintenance. Only a few short scattered sections have any remaining live markers. It is urgent that all these boundaries are attended to in this densely populated area. All internal management zone boundaries should be demarcated by ring-painting trees in the standard way. Red paint will be used to indicate Nature Reserve; yellow for 'buffer' zones. Sign boards will be erected wherever prominent footpaths cross (external and internal) boundaries.

Patrol and protection: Twelve patrol teams each comprising one Forest Guard and two patrolmen will be constituted with responsibility for safeguarding ranges as per the twelve forest stations. Men will be rotated between patrol teams and teams will be moved periodically between ranges. Patrol routes and checkpoints

will be established throughout the reserve. An incentive scheme will be instituted to reward success in curbing illegal activities.

Public access and community needs: One Forest Officer and two Forest Rangers (based at Najjembe and Maligita) will assume responsibility for community outreach programmes including the development of tourism activities, Joint Forest Management programmes within the reserve and community tree-planting programmes outside the boundary. A programme of village meetings should be instituted and developed to explain and discuss management of the reserve, and in particular the management zones as they are established. The staff will be facilitated as indicated under infrastructure.


Table 16.4: Summary Table of biodiversity values for Mabira Forest Reserve

Criteria	Trees & Shrubs	Birds	Mammals	Butterflies	Moths	Overall
Total No of species known	312	287	23	199	97	
No. of restricted range species (< 5 forests)	9	37	-	27	7	
Species unique for forest (list)	<i>Caesalpinia volkensii</i>	Tit Hylia	None	<i>Epitola catuna</i> <i>Pseudathyma plutonica</i> <i>Neptis trigonophora</i> <i>Sallya natalensis</i> <i>Acraea rogersi</i> <i>Caenides dacena</i>	<i>Orthogoniopitulum</i> sp. C.	9 spp
Uganda endemics (list)	None	none	<i>Crocidura selina</i>			4 spp
Albertine Rift endemics (list)	<i>Grewia pubescens</i>	none	None	none	None	1 spp
Species diversity (score and rank)	6.5 (26=)	6.5 (24=)	5.4 (4.0)	6.9 (25=)	5.8 (30=)	6.4(22=)
Species rarity value (Score & rank)	7.2 (29=)	6.6 (14=)	5.4 (22=)	5.6 (15=)	6.8 (+5=)	6.7(19=)
Overall biodiversity score 13						



Appendix B.5

Map of Forest Reserves and the Proposed Wayleave





Appendix B.6

Archaeological Assessment Report

**Phase 1: Archaeological Assessment for the Proposed Bujagali
Interconnection Project at Kawanda Sub- Station, Uganda.**

*By D. Kiyaga- Mulindwa and E R Kamuhangire
November 2006*

Table of Contents

Introduction.....	03
Study team.....	04
Methodology.....	05
Survey results.....	06
Discussion.....	07
References.....	08
Appendices.....	09

Introduction

An Impact Assessment of development project is a requirement of the National Environment Management Authority (NEMA) in Uganda (Ug. Govt 1995). It is also a requirement of other development funding agencies such as the African Development Bank (AfDB) and World Bank (WB 1994). This exercise is one aspect in the overall fulfillment of Impact Assessment requirement for the proposed Bujagali Interconnection project.

The main objective of this exercise was to determine if there was any possible adverse impact of project- related activities to the Archaeological resource in the direct impact zone. This survey was restricted to the area of Kawanda, in Wakiso District, which is the proposed location of the sub-station. The following is our observations and assessment in line with the main objective of the study during and after the archaeological impact assessment survey.

Study team

1. Professor D. Kiyaga Mulindwa, Professor of History, Kyambogo University, Uganda. Archaeologist, Museologist with specific Research interest in African Cultural History and several publications in these areas, especially the African Iron Age.
2. Dr. Ephraim R Kamuhangire, Commissioner Museums and Monuments of Uganda. A Historian with specific interest in Ethno- Archaeology .
3. Nelson A. Abiti, Photographer and Conservator, with specific interest in visual history and heritage management.

Methods

Methods of recognizing sites with evidence of heritage resources are varied. The general idea is to be able to recognize surface indicators of what could lie in sub surface levels to warrant further investigation through opening up test excavations. In the case of Kawanda no mapping was available and it was decided to use the Garmin GPS 76 to give the coordinates for the required relevant points. These will be down loaded to produce a map for the subsequent final version of this report.

The foot survey entailed walking the entire area since this was relatively small site. Surface or reconnaissance survey was carried out to locate archaeological features and to recover surface pottery sherds as indicator of where human activity might have taken place. This was done by walking narrow transects of 3 metres each by the three members of the team and each noting any of these indicators in his transect. Where these were noted, the location was immediately recorded and entered into the GPS.

The foot survey sought out features and artifacts, particularly potsherds. Both the physical extent and concentration of the pottery scatter are obvious indicators of the physical extend of the site as well as the concentration of human activity in that particular area.

Specific locations that seem to point to the concentration of human activity such as pottery scatter or heaps / mounds of soils or patches of ash, normally call for further investigation, such as test excavation to check on what could be lying in the sub-terrain levels . Excavation produced artifacts *in situ* and are also instrumental in displaying stratigraphic accumulation of cultural deposits which could give the chronological sequence of the occupation of that site.

A few test excavations were tried out at the proposed Kawanda sub-station for exactly the same objective as notified above.

Results.

We were directed to the proposed Kawanda – sub station by the company lawyer. As of now the site is free of occupants or any obvious permanent encumbrances.

The site situated on a hillock and estimated at about a hectare in size and is covered by rush undergrowth. To the east of the site is a patch of sweet potatoes and at the crest is another patch of sweet potatoes and some maize garden. The site is crossed by a village road, from north-west to south east.

The hill is generally covered by black top soil immersed in lateritic gravels. The site is in the middle of heavily settled area and we have a reason to believe that the settlement here has been of some antiquity since burial ground and related court of Sekabaka Sunna II (1856) at Wamala is a few kilometers west of this site. The soil composition is not the best for crop agriculture in this area and this may account for the sparse agricultural activity we noted. Furthermore, to the east of the site, we encountered recent trench which was sunk about 1.0 metres deep and equally long which exposed a bare rock. The top soil on this hillock seems to be thin in most places and underlain by bedrock, very close to the surface. This would have made settlement in earlier times which involve construction of post, mud, and wattle houses at the top of this hill, less attractive if there were easier spots lower down the slopes for such activity.

The survey revealed three house foundations one at (N00°24.577' E032°32.604') the other at (N00°24.655' E032°32.618') and the third at (N 00°24.558' E 32°32.666'). These are foundations of recent houses which were broken down during evictions. A foot survey was conducted with transects running south to north and each measuring 3 metres apart. These were walked looking out for features and artifacts. About 20 potsherds were recovered from this foot survey. These sherds were mainly from pots with rouletted decorative motifs, especially on the neck of the pot. Such decorative motifs are quite common in the Great Lakes area and in Buganda in particular and are datable from recent- to- modern. These potsherds cannot be regarded as unique archaeological finds. Four shovel test pits were sunk to test the sub- surface level for possible artifacts of antiquity at the Kawanda sub- station site and nothing was recorded.

Discussion

Kawanda sub- station is well located as it is free of any obvious encumbrances other than the few quick growing crops. Archaeologically the stratigraphy of the site has shown very light and sparse cultural deposits. From all indications even such deposits are obviously of modern times, so the area shows no historical or archaeological resource that would be threatened by the project activities. The potsherds recovered fall within what is referred to as rouletted ware. This pottery type is widely used by settled agricultural communities in this area of the interlacustrine region; it spreads up to the western province of Kenya and even the areas of northern Tanzania, south of Lake Victoria. However, in the chronological sequence, which has helped us so far to date various community migrations and settlements in this area, this pottery style is very recent. At Kawanda, what was recovered is mainly string or knotted strip roulettes, some of which are still being made and used to this day.

References

N. David and Vidal, P. 1977 “ The Nana- Mode Village site and the Prehistory of the Ubangian- speaking peoples” *West African Journal of Archaeology* 7: 17-56

D. Drost, 1967 *Topferei in Afrika: technologie* Voroffentlichungen des museums fur Volkerkunde zu heipzig 15. Berlin

D. Kiyaga- Mulindwa, 2006, “The Archaeology and Ethnography of Mutunda, A Nilotic/ Bantu Nexus in Northern Uganda”, *Studies in the African Past*, Vol.5, 224-242

R.Soper, 1985”Roulette decoration on African pottery: technical considerations, dating and distribution” *The African Archaeological Review* 3:29-51

Uganda Government, 1995, The National Environment Statue, UPPC, Entebbe

World Bank, 1994, Environment Assessment Source book, Vol.1 World Bank Washington D.C

Appendices.

Photos A



Kawanda Potsherds (Rouletted decoration)



Kawanda Potsherds (Rouletted decoration)



Kawanda foot survey



Quarry trench- Kawanda